




Flows in the River Murray System – October 2021

Flows in the River Murray System vary widely depending on a range of factors, including rainfall, inflows, evaporation, and demand for water for human use.

At any given time, water flowing through the river is destined for various uses, including irrigation, industry, communities, the environment, and meeting South Australia's flow entitlement. The exact mix of these flow components is determined by demand and water availability, amongst other factors.

The graphic below is indicative of how water flow is managed throughout the seasons across a typical year.



* including meeting South Australia's flow entitlement

Water for the environment

Overall, water for the environment is a small percentage of the total water used in the Murray-Darling Basin.

The volume of water for the environment used under water entitlements has increased slightly over the past 5 years, as more water became available. The average use over this period was 20.4% of the total water used in the Basin.

Importantly, water held for the environment uses the same entitlement framework as consumptive users. In any given year the amount of water available for delivery to key environmental sites is determined based on the same rules that apply to all other consumptive water uses.

Types of water for the environment

Water for the environment can be categorised as:


- 'Planned' environmental water water that flows through the river system that is typically managed through rules outlined in state water resource plans and is used to improve the health of the environment.
- **'Held' environmental water** is water that is held by government agencies and is delivered where and when it is needed in the river system. It is water that is allocated to relevant government agencies through water licenses for environmental use.

Who holds and manages water for the environment

(based on the long term average diversion limit for entitlements held for the environment at June 2020)

	Water manager	% of total
•	Commonwealth Environmental Water Holder Jointly held Victoria New South Wales South Australia	66% 23% 5% 4% 2%

Source: Southern Connected Basin Environmental Watering Committee Annual Report



Information in the figure above is for the month of October 2021 and may not include recent rainfall or delivery of water for the environment in the River Murray system. Information in this figure is an average estimate over the past month and formal accounts from Basin state governments may vary. Water for the environment in the figure above represents water that is held by environmental water holders, through entitlements. Other water that flows through the river can also achieve environmental outcomes.

River flow information

The October flow to South Australia comprised of water for South Australia's entitlement, traded volumes, water for the environment and unregulated flows. When rainfall falls downstream of storages and cannot be captured, it's referred to as unregulated flow. This commonly occurs when there is persistent, high inflows downstream of Hume Dam, or as the result of a spill at Hume Dam or the Menindee Lakes – these flows all significantly boost flows into the Murray. Water for the environment consisted of releases from Hume Dam (mostly Barmah–Millewa environmental water allocation) and water delivered from the Goulburn, Broken, Campaspe, Loddon and lower Darling River at weir 32.

For the latest information on water for the environment see the River Murray weekly report.

Intended environmental outcomes

Water for the environment takes time to move through the system. Water from past watering events is still moving through the River Murray as return flows. Environmental water holders can also use water for the environment by extracting allocations directly from the river. These allocations are often used for small-scale watering events rather than having water delivered from a storage.

Location	Return flows used	Intended environmental outcome(s)
River Murray delivery	n/a	 delivery of water following River Operations airspace management at Hume maintain connections between the River Murray and its forests, wetlands and creeks
Millewa Forest	n/a	provide flows through Gulpa Creek to support Australasian bitterns and large bodied native fish
Barmah Forest	n/a	 replenish refuge pools to maintain water quality, fish and turtle populations maintain connectivity between the forest and the river remove accumulated organic matter and cycle carbon in the river system to minimise the risk of hypoxic blackwater
Edward/Kolety–Wakool River system	Yes	support native fish breeding and recruitment
Lower Broken Creek	n/a	protect and increase native fish populations and avoid excessive build-up of azolla fern
Goulburn River	n/a	 protect and boost populations of native fish and turtles maintain abundant and diverse waterbugs increase water dependent plants in the river channel and banks
Campaspe River	n/a	 provide habitat to help protect and boost native fish maintain resident platypus numbers
Pyramid Creek	n/a	 maintain connectivity between pools provide habitats for native fish and waterbugs
Gunbower Forest	Yes	 fill of key wetlands and surrounding forest, supporting the growth and recruitment of wetland and river red gums vegetation increase water depth and quality to support native small-bodied fish, turtles and frogs
Gunbower Creek	Yes	maintain breeding habitat and food resources for native fish such as Murray cod
Central Murray Wetlands	Yes	create habitat for native waterbirds, frogs and turtlesstimulate the germination of aquatic vegetation
Guttrum Forest	No	 support the growth of adult river redgums and river red gum saplings promote the growth and re-establishment of aquatic vegetation and tall marsh vegetation maintain the depth of the wetland to support frogs and waterbird feeding and breeding
Mid-Murray private wetlands	No	 provide habitat for southern bellfrogs and waterbirds maintain wetland vegetation condition
Mid-Murray ephemeral creeks (Tuppal, Cockrans–Jimaringle, Murrain–Yarrien, Yarrien, Thule)	No	 provide habitat for southern bellfrogs and waterbirds maintain fringing river redgum health
The Pollack (Koondrook– Perricoota)	No	 provide habitat for small bodied native fish and waterbirds maintain wetland vegetation condition
Little Forest (Koondrook– Perricoota)	No	 provide habitat for waterbirds maintain wetland vegetation condition
Lindsay, Mulcra and Wallpolla islands (Lake Wallawalla and Wallpolla Horseshoe)	No	 provide shallow-water, open-water and shoreline habitat provide foraging habitat and breeding opportunities for frogs, yabbies, waterbirds and turtles
Hattah lakes	No	 support the growth and recruitment of wetland and river red gums vegetation inundate dry wetlands to release nutrients to increase food web productivity provide waterbird breeding habitat and support small-bodied fish
Lower Darling River (Baaka)	n/a	 small increase to baseflows to improve the condition of the Lower Darling (Baaka) assist native fish population recovery by supporting breeding opportunities of primarily Murray cod (and potentially golden perch and silver perch) this spring
Great Darling Anabranch	No	 reconnection event allowing a "Golden Highway" for a good number of juvenile golden perch to travel from the Menindee Lakes system and into the Murray River
Lower Murray wetlands	n/a	 wet fringing vegetation communities to improve their condition provide a range of open water, shallow water and inundated lignum habitats for waterbird feeding and breeding

More information on river flows and water for the environment

Live River Data riverdata.mdba.gov.au River Murray Weekly Report mdba.gov.au/river-information/weekly-reports Water sharing in the River Murray www.mdba.gov.au/river-information/water-sharing Water use in catchments www.environment.gov.au/water/cewo/catchment FLOW Monitoring, Evaluation and Research flow-mer.org.au

Delivering water for the environment mdba.gov.au/managing-water/water-for-environment/water-over-time