




Macquarie-Castlereagh Long Term Water Plan

Part A



© 2018 State of NSW and Office of Environment and Heritage

With the exception of photographs, the State of NSW and Office of Environment and Heritage are pleased to allow this material to be reproduced in whole or in part for educational and non-commercial use, provided the meaning is unchanged and its source, publisher and authorship are acknowledged. Specific permission is required for the reproduction of photographs.

The Office of Environment and Heritage (OEH) has compiled this report in good faith, exercising all due care and attention. No representation is made about the accuracy, completeness or suitability of the information in this publication for any particular purpose. OEH shall not be liable for any damage which may occur to any person or organisation taking action or not on the basis of this publication. Readers should seek appropriate advice when applying the information to their specific needs.

All content in this publication is owned by OEH and is protected by Crown Copyright, unless credited otherwise. It is licensed under the [Creative Commons Attribution 4.0 International \(CC BY 4.0\)](#), subject to the exemptions contained in the licence. The legal code for the licence is available at [Creative Commons](#).

OEH asserts the right to be attributed as author of the original material in the following manner: © State of New South Wales and Office of Environment and Heritage 2018.

Cover photo: Macquarie Marshes, John Spencer OEH

Published by:

Office of Environment and Heritage
59 Goulburn Street, Sydney NSW 2000
PO Box A290, Sydney South NSW 1232
Phone: +61 2 9995 5000 (switchboard)
Phone: 131 555 (environment information and publications requests)
Phone: 1300 361 967 (national parks, general environmental enquiries, and publications requests)
Fax: +61 2 9995 5999
TTY users: phone 133 677, then ask for 131 555
Speak and listen users: phone 1300 555 727, then ask for 131 555
Email: info@environment.nsw.gov.au
Website: www.environment.nsw.gov.au

Report pollution and environmental incidents

Environment Line: 131 555 (NSW only) or info@environment.nsw.gov.au

See also www.environment.nsw.gov.au

ISBN 978 1 76039 978 8

OEH 2018??/0696

Month 2018

Find out more about your environment at:

www.environment.nsw.gov.au

Contents

Acknowledgement of Traditional Owners	vii
Abbreviations	viii
Glossary	ix
Summary	xiv
1. Introduction	16
1.1 Approach to developing the LTWP	16
1.2 Implementing the Macquarie-Castlereagh LTWP	17
1.3 LTWP document structure	17
1.4 Planning units	18
2. Environmental assets	20
2.1 Audit of water-dependent (priority) environmental assets	20
3. Ecological objectives and targets	22
3.1 Native fish objectives	22
3.2 Native vegetation objectives	25
3.3 Waterbird objectives	28
3.4 Priority ecosystem function objectives	31
3.5 Objectives for other species (flow-dependent frogs)	36
4. Environmental water requirements	38
4.1 Developing EWRs	38
4.2 Flow threshold estimates for flow components in the regulated Macquarie-Castlereagh	41
4.3 Catchment-scale EWRs for Macquarie-Castlereagh	44
4.4 Important flow regime characteristics needed to deliver LTWP objectives	51
4.5 Changes to the flow regime	59
5. Risks, constraints and strategies	66
5.1 Risks and constraints to meeting EWRs in the Macquarie-Castlereagh WRPA	67
5.2 Non-flow related risks and constraints to meeting LTWP objectives	70
5.3 Dealing with the risk of climate change	76
6. Water management under different water availability scenarios	78
6.1 Prioritisation of ecological objectives and watering in regulated river reaches	78
6.2 Protection of ecologically important flow components in unregulated river reaches	85
7. Going forward	86

7.1	Potential further actions for water management.	86
7.2	Cooperative water use	86
7.3	Complementary actions	87
7.4	Measuring progress	92
7.5	Review and update	93
References	94	
Appendix A	LTWP objectives relevant to each planning unit in the Macquarie-Castlereagh WRPA	98
Appendix B	RAS guidelines	100

List of tables

Table 1	Native fish ecological objectives	23
Table 2	Native vegetation ecological objectives	27
Table 3	Waterbird ecological objectives	30
Table 4	Ecosystem function objectives	33
Table 5	Objectives for other species (flow-dependent frogs)	37
Table 6	Description of function role provided by flow components	39
Table 7	Flow threshold estimates for flow components in the regulated Macquarie-Castlereagh WRPA	41
Table 8	Catchment-scale EWRs for Macquarie-Castlereagh WRPA	44
Table 9	Important flow regime characteristics needed to deliver LTWP objectives	51
Table 10	Predicted inundated areas for specific flow scenarios	61
Table 11	Water management related risks and constraints to meeting and potential strategies for managing them	67
Table 12	Non-flow management related risks and constraints to meeting LTWP ecological objectives	71
Table 13	Projected change in rainfall patterns and weather in the Central West and Orana district from 2020 to 2079 due to climate change NARCLiM (nd)	76
Table 14	Priority LTWP objectives and flow components in a very RAS	79
Table 15	Priority LTWP objectives and flow components in a dry RAS	80
Table 16	Priority objectives and flow components in a moderate RAS	81
Table 17	Priority LTWP objectives and flow components in a wet RAS	83
Table 18	Potential management strategies for meeting LTWP objectives in unregulated river reaches of the Macquarie Castlereagh WRPA	85
Table 19	Recommended further investment and projects to improve environmental water outcomes in the Macquarie-Castlereagh WRPA	88
Table 20	Default matrix for determining the RAS (MDBA 2012b)	100

List of figures

Figure 1	Aerial view of the Macquarie Marshes. Photo John Spencer/OEH	vii
Figure 2	The Macquarie-Castlereagh WRPA showing the division of planning units	19
Figure 3	The five criteria for the identification of environmental assets	20
Figure 4	Murray cod. Photo Gunther Schmida	25
Figure 5	Intermediate egret in breeding plumage. Photo John Spencer OEH	29
Figure 6	Salmon striped frog. Photo: Joanne Ocock/OEH	36
Figure 7	A simplified conceptual model of the role of flow regime components	39
Figure 8	Schematic of the main watercourses and streamflow gauges used for EWRs (Not to scale)	43
Figure 9	Burrendong Dam offtake tower with cold water curtain. Photo: Tim Hosking/OEH	59
Figure 10	Areas of the inundation in the Macquarie Marshes based on flow scenarios of different sizes listed in Table 10 from Thomas et al. (in prep)	62
Figure 11	Occurrence of events in Macquarie Marshes, based on flow scenarios of different sizes listed in Table 10 from Thomas et al. (in prep).	63
Figure 12	Sunset over Macquarie Marshes. Photo John Spencer/OEH	66
Figure 13	Golden perch. Photo: Gunther Schmida	75
Figure 14	Burrendong Dam offtake tower with cold water curtain. Photo: Tim Hosking/OEH	87

Acknowledgement of Traditional Owners

The Office of Environment and Heritage pays its respect to the Traditional Owners and their Nations of the Murray-Darling Basin. The contributions of earlier generations, including the Elders, who have fought for their rights in natural resource management are valued and respected.

In relation to the Macquarie-Castlereagh Water Resource Plan Area, the Office of Environment and Heritage pays its respects to the Traditional Owners – the Gomeroi/Kamilaroi, Ngemba, Ngiyampaa, Wailwan, Wiradjuri Nations – past, present and future. We look forward to building upon existing relationships to improve the health of our rivers, wetlands and floodplains including in recognition of their traditional and ongoing cultural and spiritual significance.



Figure 1 Aerial view of the Macquarie Marshes. Photo John Spencer/OEH

Formatted: None, Space Before: 6 pt, After: 6 pt, Line spacing: single, Don't keep with next, Don't keep lines together

Formatted: None, Space Before: 6 pt, After: 6 pt, Line spacing: single, Don't keep with next, Don't keep lines together

Abbreviations

Basin Plan	Murray-Darling Basin Plan
BWS	Basin-wide environmental watering strategy
CAG	Customer Advisory Group
CAMBA	China-Australia Migratory Bird Agreement
CEWH	Commonwealth Environmental Water Holder
CEWO	Commonwealth Environmental Water Office
DO	Dissolved oxygen
DOC	Dissolved organic carbon
DOI – Water	NSW Department of Industry–Water
EWA	Environmental water allowance
EFRG	Environmental Flows Reference Group (the Macquarie Cudgegong EWAG)
EWAG	Environmental Water Advisory Group
EW	Environmental water requirement
HEW	Held environmental water
JAMBA	Japan-Australia Migratory Bird Agreement
LLS	NSW Local Land Services
LTWP	Long Term Water Plan
MDBA	Murray-Darling Basin Authority
MER	Monitoring, evaluation and reporting
mg/L	milligrams per litre
ML	megalitre
ML/d	megalitres per day
m/s	metres per second
NPWS	NSW National Parks and Wildlife Services
NRAR	NSW Natural Resources Access Regulator
NSW	New South Wales
OE	NSW Office of Environment and Heritage
PEW	Planned environmental water
ROKAMBA	Republic of Korea-Australia Migratory Bird Agreement
SDL	Sustainable diversion limit
WRP	Water resource plan
WRPA	Water resource plan area
WSP	Water sharing plan

Glossary

Actively managed floodplain	The floodplains and wetlands that can be inundated by flows from regulated rivers (see 'Regulated river').
Adaptive management	A procedure for implementing management while learning about which management actions are most effective at achieving specified objectives.
Allocation	The volume of water made available to a water access licence or environmental water allowances in a given period by DOI-W, which is typically determined within the context of demand, inflows, and stored water.
Alluvial	Comprised of material deposited by water.
Bankfull flow	River flows at maximum channel capacity with little overflow to adjacent floodplains. Engages the riparian zone, anabranches and flood runners and wetlands located within the meander train. Inundates all in channel habitats including all benches, snags and backwaters.
Baseflow	Reliable background flow levels within a river channel that are generally maintained by seepage from groundwater storage, but also by surface inflows. Typically inundates pools and riffle areas.
Basin Plan	The Basin Plan as developed by the Murray-Darling Basin Authority under the <i>Water Act 2007</i> .
Biota	The organisms that occupy a geographic region.
Blackwater	Occurs when water moves across the floodplain and releases organic carbon from the soil and leaf litter. The water takes on a tea colour as tannins and other carbon compounds are released from the decaying leaf litter. The movement of blackwater plays an important role in transferring essential nutrients from wetlands into rivers and vice versa. Blackwater carries organic carbon compounds which are the basic building block of the aquatic food web and an essential part of a healthy river system. See 'hypoxic blackwater' below for those occasions this can cause significant water quality issues.
Carryover	Water allocated to water licences or environmental water accounts that remains unused at the end of the water year which, under specific circumstances, may be held over and used in the following water year/s.
Cease-to-flow	The absence of flowing water in a river channel. Partial or total drying of the river channel. Streams contract to a series of isolated pools.
Cease-to-pump (access rule in WSP)	<p>This is a low flow restriction on access to water for Works Approval Licences. Generally, licenced take is not permitted:</p> <ol style="list-style-type: none"> 1. from in-channel pools and from natural off-river pools when the water level is lower than its full capacity 2. from pump sites when there is no visible flow <p>These rules typically apply unless there is a commence-to-pump access rule that specifies a higher flow rate that licence holders can begin pumping.</p>
Constraints	The physical or operational constraints that effect the ability to meet flow levels in the river or wetlands. These include limitations on the delivery of water from storages to extraction or diversion points or to wetland areas. Constraints may include structures such as bridges that can be affected by higher flows, or the volume of water that can be carried through the river channel, or scheduling of downstream water deliveries from storage.
Consumptive water	Water that is removed from surface and groundwater systems without return to a water resource system (such as water removed from a river for agriculture, mining or urban use).

Cultural water-dependent asset	A place that has social, spiritual and cultural value based on its cultural significance to Aboriginal people and is reliant upon surface or groundwater supply for its values.
Cultural water-dependent value	An object, plant, animal, spiritual connection or use that is dependent on water and has value based on its cultural significance to Aboriginal people.
Discharge	The amount of water moving through a river system, most commonly expressed in megalitres per day (ML/d).
Dissolved Organic Carbon (DOC)	A measurement of the amount of carbon from organic matter that is soluble in water. DOC is transported by water from floodplains to river systems and is a basic building block available to bacteria and algae that are food for microscopic animals that are in turn consumed by fish larvae, small bodied fish species, yabbies and shrimp. DOC is essential for building the primary food webs in rivers and ultimately generates a food source for large-bodied fish like Murray cod and golden perch and predators such as waterbirds.
Ecological function	The resources and services that sustain human, plant and animal communities and are provided by the processes and interactions occurring within and between ecosystems.
Ecological objective	The defined goal for a state, condition or characteristic of an ecological asset or function.
Ecological target	Level of measured performance that must be met to achieve the defined objective. The targets in this long term water plan are SMART (Specific/Measurable/Achievable/Realistic/Time-bound).
Ecological value	An object, plant or animal which has value based on ecological significance.
Ecosystem	A biological community of interacting organisms and their physical environment. It includes all the living things in that community, interacting with their non-living environment (weather, earth, sun, soil, climate and atmosphere) and with each other.
Environmental asset	An object, area, species or ecological community that has ecological value, e.g. wetlands, species, riverine forests.
Environmental Water Allowance (EWA)	An allowance of water provided by the <i>Water Sharing Plan Water Sharing Plan for the Macquarie and Cudgegong Regulated Rivers Water Source</i> for environmental purposes.
Environmental water	Water for meeting the requirements of water-dependent ecosystems. It provides a multitude of benefits to not only the environment, but to communities, industry and society. It includes held environmental water and planned environmental water.
Environmental water requirement (EWR)	The water required to support the completion of all elements of a lifecycle of an organism or group of organisms (taxonomic or spatial), consistent with the objective/target, measured at the most appropriate gauge. Includes all water in the system including natural inflows, held environmental water and planned environmental water.
Five-year rolling period	Period of time used for monitoring and evaluation. This looks at every 5-year period on a continuous basis.
Flow component	The type of flow in a waterway defined by its magnitude, season, shape and role (e.g. bankfull, spring fresh for native fish breeding).
Flow-dependent	In the context of this plan, an ecosystem, community or species that depends on periodic or sustained inundation, waterlogging or significant inputs of surface water for part or all of its lifecycle. See also 'water-dependent.'
Flow-dependent frog	Frog species that respond with breeding behaviour to wetland filling from surface water flows. Other species respond primarily to local rainfall events but may also utilise wetlands if rainfall coincides with filling.
Flow regime	The pattern of flows in a waterway or wetland over time that will influence the response and persistence of plants, animals and their ecosystems.

Freshes	Temporary in-channel flow pulse that typically happens in response to rainfall or release from water storages. Very important for a range of ecological values.
General security	A category of water entitlement where water is made available to licence holder accounts based upon dam inflows. This forms the majority, by volume, of water entitlements in the regulated sections of the river including irrigation. The Commonwealth and NSW governments hold general security access licences to use for the environment.
Groundwater	Water that is located below the earth's surface in soil pore spaces and in the fractures of rock formations. Groundwater is recharged from, and can eventually flow to, the surface naturally.
Held environmental water	Water available under a water access right, a water delivery right, or an irrigation right for the purposes of achieving environmental outcomes (including water that is specified in a water access right to be for environmental use).
Hydrological connectivity	The link of natural aquatic environments.
Hydrology	The occurrence, distribution and movement of water.
Hypoxic blackwater	<p>Occurs when dissolved oxygen (DO) levels fall below the level needed to sustain native fish and other water dependent species. Bacteria feed on dissolved organic carbon use oxygen in the water. When they multiply rapidly their rate of oxygen consumption can exceed the rate at which oxygen can be dissolved in the water and oxygen levels fall and a hypoxic (low oxygen) condition occurs. This is typically not a frequent problem in the Macquarie-Castlereagh WRPA.</p> <p>Dissolved oxygen is measured in milligrams per litre (mg/litre). Generally native fish begin to stress when DO levels fall below 4 mg/litre. Fish mortality occurs when DO levels are less than 2 mg/litre.</p>
Key environmental asset	An asset that is identified for its special conservation significance based on selected temporal and spatial criteria.
Key ecological value	A species or community that is identified for its special conservation significance based on selected temporal and spatial criteria. Examples include Murray cod or river red gum woodlands.
Large fresh	A high-magnitude flow pulse that remains in-channel. May engage flood runners with the main channel and inundate low-lying wetlands. Connects most in channel habitats and provides partial longitudinal connectivity, as some low-level weirs and other in channel barriers may be drowned out. Highly important for aquatic ecosystems.
Lateral connectivity	The flow linking rivers channels and the floodplain
Long Term Water Plan (LTWP)	Plans required of Basin States by the Murray-Darling Basin Plan. Long term water plans give effect to the <i>Basin-wide Environmental Watering Strategy</i> relevant for each river system and will guide the management of water over the longer term. These plans will identify the environmental assets that are dependent on water for their persistence, and match that need to the water available to be managed for or delivered to them. The plan will set objectives, targets and watering requirements for key plants, waterbirds, fish and ecosystem functions. OEH is responsible for the development of nine plans for river catchments across NSW, with objectives for five, 10 and 20-year timeframes.
Longitudinal connectivity	The flow link along the length of a river.
Overbank flow	Flows that spill over the riverbank or extend to floodplain surface flows.
Planned environmental water	Water that is committed by the Basin Plan, a water resource plan or a plan made under state water management law to specifically achieve environmental outcomes.

Planning Unit (PU)	A geographical division of a water resource plan area based on water requirements (in catchment areas in which water is actively managed), or a sub-catchment boundary (all other areas).
Population structure (ecological)	A healthy population structure has individuals in a range of age and size classes. These populations demonstrate regular recruitment, good numbers of sexually mature individuals and genetic variation.
Priority ecological function	In the context of this plan, is a water-dependent ecological function that can be influenced by environmental water.
Priority environmental asset	In the context of this plan, is a place of particular ecological significance that contains values and functions that are water-dependent and can be influenced by environmental water.
Ramsar Convention	An international treaty to maintain the ecological character of key wetlands.
Recruitment (ecological)	Successful development and growth of offspring within a population; such that they can contribute to the next generation.
Refuge (ecological)	An area which provides conditions to assist individuals within a population of plants or animals to survive through a period of decreased water availability.
Registered cultural asset	A cultural water-dependent asset that is registered in the NSW Aboriginal Heritage Information Management System.
Regulated river	A river that is gazetted under the <i>NSW Water Management Act 2000</i> . Flow is largely controlled by major dams, water storages and weirs. River regulation brings more reliability to water supplies but has interrupted the natural flow characteristics and regimes required by native fish and other plant and animal to breed, feed and grow.
Riffle	A rocky or shallow part of a river where river flow is rapid and broken.
Riparian	The part of the landscape adjoining rivers and streams that has a direct influence on the water and aquatic ecosystems within them.
Small fresh	Low-magnitude in-channel flow pulse. Unlikely to drown out any significant barriers, but can provide limited connectivity and a biological trigger for animal movement.
Supplementary access licence	A category of water entitlement where water is made available to licence holders during periods of high river flows, subject to triggers. Water can be taken and debited from licence accounts during a declared period. The Commonwealth and NSW governments hold supplementary access licences to use for the environmental.
Surface water	Water that exists above the ground in rivers, streams creeks, lakes and reservoirs. Although separate from groundwater, they are interrelated and over extraction of either will impact on the other.
Sustainable diversion limit (SDL)	The total amount of water that can be extracted from Murray-Darling Basin rivers for human uses. Water in the system above the SDL is protected to achieve environmental outcomes.
Thermal (including coldwater) pollution	The artificial lowering or raising of water temperature that occurs downstream of large dams, particularly during warmer months when they stratify.
Unregulated river	A waterway where flow is mostly uncontrolled by dams, weirs or other structures.
Very low flow	Small flow in the very-low flow class that joins river pools, thus providing partial or complete connectivity in a reach. Can improve DO saturation and reduce stratification in pools.
Water quality management plan (WQSP)	A document prepared by state authorities and accredited by the Commonwealth under the Basin Plan. Makes up part of the Water Resource Plan. It aims to provide a framework to protect, enhance and restore water quality in each water resource plan area.

Water resource plan (WRP)	A document prepared by state authorities and accredited by the Commonwealth under the Basin Plan. The document describes how water will be managed and shared between users in an area.
Water resource plan area (WRPA)	Catchment-based divisions of the Murray-Darling Basin defined by a water resource plan.
Water sharing plan (WSP)	A plan made under the NSW <i>Water Management Act 2000</i> that sets out specific rules for sharing and trading water between the various water users and the environment in a specified water management area. A water sharing plan will be a component of a water resource plan.
Water-dependent	A term used in the Basin Plan. In the context of this plan, an ecosystem, community or species that depends on periodic or sustained inundation, waterlogging or significant inputs of surface water for part or all of its lifecycle.

Summary

Rivers, creeks, wetlands and floodplains play a vital role in sustaining healthy communities and economies. They provide productivity and connections across the landscape for people, plants and animals with benefits that extend well beyond the river bank.

Over the past 200 years, many rivers, wetlands and floodplains in New South Wales (NSW) have had their natural flow regimes disrupted because of dams, weirs, floodplain development, and water regulation and extraction.

The Macquarie-Castlereagh Long Term Water Plan (LTWP) is an important step to describing the flow regimes that are required to maintain or improve environmental outcomes in the Macquarie-Castlereagh catchment. The plan identifies water management strategies for maintaining and improving the long-term health of the Macquarie-Castlereagh's riverine and floodplain environmental assets and the ecological functions they perform. This includes detailed descriptions of ecologically important river flows and risks to water for the environment.

Importantly, the LTWP does not prescribe how environmental water should be managed in the future, rather it will help water managers and advisory groups, such as the Macquarie-Castlereagh Environmental Flows Reference Group, make decisions about where, when and how water can be used to achieve agreed long-term ecological objectives.

The LTWP looks at all sources of water and how these can be managed to help support environmental outcomes in the catchment.

Background to Long Term Water Plans

The Basin Plan (Pt 4, Ch. 8) establishes a framework for managing environmental water at the Basin and catchment-scale. The framework is designed to ensure environmental water managers work collaboratively to prioritise water use to meet the long-term needs of native fish, water-dependent native vegetation and waterbirds and co-ordinate water use across multiple catchments to achieve Basin-scale outcomes.

The *Basin-wide Environmental Watering Strategy* (BWS) and LTWPs are central features of this framework. The BWS establishes long-term environmental outcomes and targets for the Basin and its catchments. LTWPs, which apply to water resource plan areas (WRPAs) (catchment-scale), must contribute to the achievement of the BWS by identifying:

- priority environmental assets and functions in a WRPA
- ecological objectives and ecological targets for those assets and functions
- environmental watering requirements needed to meet those targets and achieve the objectives.

Water resource plans (WRPs) must have regard to LTWPs.

The Macquarie-Castlereagh Long Term Water Plan

The Macquarie-Castlereagh LTWP is one of nine plans being developed by the Office of Environment and Heritage (OEH) to cover the NSW portion of the Murray-Darling Basin. Development of the LTWP has involved six main steps.

- Undertaking a comprehensive **stocktake** of water-dependent environmental assets and ecosystem functions across the catchment to identify native fish, water-dependent bird and vegetation species, and river processes that underpin a healthy river system.
- Determining specific and quantifiable **objectives and targets** for the key species and functions in the Macquarie-Castlereagh WRPA.
- Determining the **environmental water requirements** (EWRs) (including volume, frequency, timing and duration) needed to achieve the objectives and targets.
- Identifying the **risks and constraints** to meeting the long-term water requirements of priority environmental assets and ecosystem functions.

- Identifying potential **management strategies** for guiding water management decisions and investment into the future.
- Identifying **complementary investments** to address risks and constraints to meeting the long-term water requirements of priority environmental assets and ecosystem functions.

The LTWP presents this information in nine chapters in two parts, with accompanying appendices.

Environmental values of the Macquarie-Castlereagh catchment

The Macquarie-Castlereagh catchment supports a range of water-dependent ecosystems, including instream aquatic habitats, riparian forests, and floodplain watercourses, woodlands and wetlands. Notably, the Macquarie Marshes are on the List of Wetlands of International Importance (the Ramsar List), and the Directory of Important Wetlands in Australia. These ecosystems benefit many water-dependent species, including threatened ecological communities, threatened, endangered and migratory waterbirds, and threatened native fish species, by providing habitat and food resources.

The ecological condition of the Macquarie-Castlereagh's water-dependent environmental assets is largely driven by flows that connect the instream benches, anabranches, floodplains and wetlands. Flows that provide these connections support organic carbon transfer and nutrient cycling, trigger movement and breeding of native fish and waterbirds, and directly impact vegetation condition and habitat availability.

Extensive local and scientific knowledge about the Macquarie-Castlereagh's riverine environmental assets and ecosystem functions underpins this LTWP. This has been collected in partnership with water managers, natural resource managers, environmental water holders, landholders, and community members. Information about the Macquarie-Castlereagh's environmental values closely aligns with material in the NSW Department of Industry *Macquarie-Castlereagh's Water Resource Plan Risk Assessment* (NSW DOI-W, in prep.).

Water for the environment

The Macquarie-Castlereagh LTWP contains ecological objectives and targets for priority environmental assets and ecosystem functions in the Macquarie-Castlereagh catchment. Priorities are defined by the Basin Plan as those assets and functions that can be managed with environmental water. The objectives and targets have been identified in this LTWP for native fish, native vegetation, waterbirds, river connectivity and flow-dependent frogs.

The objectives of this plan reflect the current understanding of outcomes that might be expected from implementation of the Basin Plan in the rivers, wetlands, floodplains, and watercourses of the Macquarie-Castlereagh catchment. The targets for each ecological objective provide a transparent means of evaluating progress towards their achievement and the long-term success of management strategies.

1. Introduction

A major catchment in the Murray-Darling Basin, the Macquarie-Castlereagh WRP is located in central-western NSW. It extends west from the Blue Mountains in the Great Dividing Range to the confluence of the Barwon River east of Brewarrina. Major towns and cities supported by the Macquarie River include Bathurst, Wellington, Dubbo and Warren. Other major rivers in the system include the Castlereagh and Bogan.

Home to the Macquarie Marshes, this catchment supports one of the largest remaining semi-permanent wetland systems in inland Australia. Listed under the Ramsar Convention, the Macquarie Marshes support a variety of important flow-dependent vegetation communities and is one of the largest colonial waterbird breeding sites in Australia.

The waterways and aquifers of the Macquarie-Castlereagh are important water resources for agricultural businesses and urban communities, with the Macquarie Marshes supporting a highly productive grazing industry.

The region supports a diverse range of flow-dependent threatened and iconic species including:

- native fish (e.g. trout cod, Murray cod, silver perch, southern purple-spotted gudgeon and eel-tailed catfish)
- vegetation communities (e.g. coolibah, black box, river red gum, lignum, reedbeds)
- waterbirds (e.g. straw-necked ibis, intermediate egret, great egret, Australian painted snipe, magpie goose, glossy ibis).

River flows in the Macquarie-Castlereagh WRP have been altered by two major headwater storages—Burrendong and Windamere dams—as well as weirs and floodplain development to support primary production. Flow volumes, as well as the regularity of small to moderate-sized flow events, have reduced compared to natural conditions as a result. These changes change the ability of the river system to support the catchment's riverine and floodplain ecosystems.

The NSW Government has developed this plan with the aim of protecting and improving the health of the Macquarie-Castlereagh's riverine and floodplain ecosystems. The plan provides information on the catchment's long-term environmental water requirements (EWRs). It also recognises the Macquarie-Castlereagh's connection and role within the Murray-Darling Basin in supporting the environmental health of downstream catchments, particularly the Barwon-Darling River.

1.1 Approach to developing the LTWP

The Macquarie-Castlereagh LTWP ('the plan') applies to the Macquarie-Castlereagh WRP and is one of nine catchment-based plans covering the NSW portion of the Murray-Darling Basin. This plan is consistent with the requirements of the Basin Plan (MDBA 2012a).

This plan is the product of best available information and engagement with water managers, natural resource managers, environmental water holders and community members. It draws together local, traditional and scientific knowledge to identify the catchment's priority environmental assets and ecosystem functions to guide the management of water to protect and restore condition over the long-term.

Development of the Macquarie-Castlereagh plan has involved six main steps.

1. **System Audit:** undertaking a comprehensive stocktake of water-dependent environmental assets and ecosystem functions across the Macquarie-Castlereagh WRP to identify native fish, water-dependent bird and vegetation species, and river processes that underpin a healthy river system.
2. **Objectives and targets:** determining specific and quantifiable objectives and targets for the key species and functions in the Macquarie-Castlereagh WRP.

3. **EWRs:** using available science and management experience to determine the water requirements (including volume, frequency, timing and duration) needed to sustain and improve the health and/or extent of priority environmental assets and ecosystem functions.
4. **Risk Assessment:** identifying the risks and constraints to meeting the long-term water requirements of priority environmental assets and ecosystem functions.
5. **Management Strategies:** identifying potential management strategies for guiding water management decisions and investment into the future.
6. **Complementary measures:** identifying actions other than water management to address risks and constraints to meeting the plan's objectives.

1.2 Implementing the Macquarie-Castlereagh LTWP

Implementation of the plan requires strong partnerships and coordination between land and water managers. This plan provides the foundation to support future coordination efforts by:

- informing annual and longer-term water management planning by environmental water managers, including strengthening collaboration between environmental water holders
- informing water planning processes that influence river and wetland health outcomes, including water sharing plans and water resource plans
- identifying opportunities for more strategic river operations by Water NSW
- helping target investment priorities for complementary actions that will effectively contribute to progressing the outcomes sought by this plan
- informing monitoring, evaluation and reporting processes for water management
- assisting to build community understanding of river and wetland health issues and management actions.

1.3 LTWP document structure

This plan is presented in nine chapters with accompanying appendices. It is divided into Part A and Part B.

Part A

- **Chapter 1** explains the background and purpose of the plan.
- **Chapters 2 and 3** identify the Macquarie-Castlereagh's water-dependent environmental assets and ecosystem functions, and articulate the environmental outcomes that are expected from implementation of the plan through ecological objectives and targets.
- **Chapter 4** provides the EWRs that are needed to support the achievement of ecological objectives over the next five, 10 and 20 years.
- **Chapter 5** describes the long-term risks and operational constraints to achieving the EWRs and ecological objectives as well as recommending management strategies to address these risks and constraints.
- **Chapter 6** identifies opportunities for the use of held and planned environmental water, and other system flows to support flow regimes to meet the EWRs of the Macquarie-Castlereagh environmental assets and values under dry, moderate and wet water resource availability scenarios.
- **Chapter 7** describes potential cooperative arrangements between government agencies and private landholders and prioritised investment opportunities to achieve the environmental outcomes described.

Part B – planning unit information

- Present the LTWP at the planning unit (PU) scale. This includes a summary of the environmental values the PU supports, and an evaluation of the impact of water resource development on local hydrology.

1.4 Planning units

The PUs shown in Figure 2 are referred to in most chapters. The PU boundaries typically align with water source boundaries in the *Macquarie-Castlereagh Water Resource Plan* (NSW Government 2018). However, some of these water sources have been amalgamated or split depending how water management for the environment can be implemented. Where there are similarities between water sources they have been amalgamated, where there are differences they have been split. It is important to note that some PUs are regulated water sources or affected by regulated water (PU 1–11). Others are unregulated and not able to be influenced by regulated water deliveries (PU 12–28).




Figure 2 The Macquarie-Castlereagh WRPA showing the division of planning units

2. Environmental assets

The Macquarie-Castlereagh WSPA supports a variety of water-dependent ecosystems, including instream aquatic habitats, floodplain woodlands and wetlands. These features are spread throughout the catchment and each has their own water requirements depending on the plants and animal species they support and ecosystem functions they perform.

2.1 Audit of water-dependent (priority) environmental assets

Schedule 8 of the Basin Plan outlines criteria for identifying water-dependent ecosystems that should be recognised as environmental assets in the Murray-Darling Basin. The criteria are designed to identify water-dependent ecosystems that are internationally important, natural or near-natural, provide vital habitat for native water-dependent biota, and/or can support threatened species, threatened ecological communities or significant biodiversity.

The Macquarie-Castlereagh's water-dependent ecosystems, which are comprised of waterbodies and surrounding water-dependent vegetation, were assessed against the Schedule 8 criteria. Significant Aboriginal cultural water-dependent sites that are registered in the Aboriginal Heritage Information Management System were also included as water-dependent assets in the LTWP. This includes areas such as recorded Aboriginal ceremony and dreaming sites, fish traps, scar trees, and water holes throughout the Macquarie-Castlereagh WSPA.

Results of the analysis are presented in Figure 3. For more information on this assessment process, refer to *Long Term Water Plan Background Information* (NSW OEH 2018).




Figure 3 The five criteria for the identification of environmental assets

Priority environmental assets in LTWP's are the assets that have been identified using Schedule 8 criteria that can be managed with planned and held environmental water. These include areas:

- subject to discretionary environmental water management, such as the Macquarie Marshes and along regulated streams, including the Macquarie River
- downstream of the regulated system that are also subject to managed environmental flows, such as the Lower Macquarie River (downstream of the Marshes) and the Lower Marthaguy Creek
- supported by implementation of the water sharing plan rules, including regulated and unregulated streams and floodplains.

Priority environmental assets may be a reach of river channel and its floodplain features at a geographic location, or a wetland complex or anabranch.

3. Ecological objectives and targets

Ecological objectives and targets have been established for priority environmental assets in the Macquarie-Castlereagh WRP (sections 3.1–3.5). They are grouped into five themes—native fish, native vegetation, waterbirds, ecosystem functions and other species. The first four themes are consistent with the BWS (MDBA 2014a). Each theme is a good indicator of river system health and is responsive to flow. The water requirements of foundational species, communities or ecosystem functions within each theme are also broadly representative of those needed by many of catchment's water-dependent species that do not fit within the five themes, such as turtles, platypus, mussels and yabbies.

The ecological objectives express the environmental outcomes that are expected from implementation of the LTWP. Their achievement will also contribute to the landscape and basin-scale environmental outcomes sought by the BWS.

The five, 10 and 20-year targets for each ecological objective provide a transparent means for evaluating progress over time and test the LTWP's management strategies and their implementation. It is hoped that these targets, if achieved, will be a suitable indicator of overall trends in ecosystem health. It is recognised that prevailing conditions such as drought and floods will influence our ability to meet targets in 5-year increments, and these will need to be taken into account in analysis of outcomes. Failure to meet targets should trigger re-assessment of the related flow regime and whether the LTWP is being implemented as intended to determine if changes are needed.

The ecological objectives for the Macquarie-Castlereagh priority environmental assets as they relate to individual PUs are listed in Appendix A. The selection of ecological objectives recognises the values that the priority environmental asset supports (e.g. native fish species, native vegetation communities, waterbirds) or the ecosystem function it performs (e.g. provides vital instream habitat).

3.1 Native fish objectives

The native fish community in the Macquarie-Castlereagh WRP consists of 19 native species recorded or expected to occur in the catchment (NSW DPI 2015). This includes seven native fish species listed as threatened or vulnerable under the *NSW Fisheries Management Act 1994* and *Environment Protection and Biodiversity Conservation Act 1999*. These species include: Darling River snail, flathead galaxias, Macquarie perch, Murray cod, purple-spotted gudgeon, silver perch and trout cod. Endangered fish populations in the Macquarie-Castlereagh include the western population of the olive perchlet and the Murray-Darling Basin population of the eel-tailed catfish.

Overall, the health of the fish community in the Macquarie-Castlereagh was rated as poor by NSW DPI Fisheries in 2015 (NSW DPI 2015). However, they found that some reaches were in moderate condition (the Bogan River and Castlereagh River systems). Reaches that were rated as being in very poor health for fish were mostly restricted to the upper catchment.

Factors that have contributed to the poor health of native fish communities in the Macquarie-Castlereagh (from NSW DPI 2015) include:

- barriers to fish movement (238 major instream structures)
- changes to flow regimes due to regulation
- degradation of in-stream habitat and riparian vegetation due to sedimentation, nutrients and de-snagging
- inappropriate land management
- the impact of introduced fish species like common carp, redfin and gambusia

- loss of large numbers of young fish through entrainment of fish in pumps and the crushing of fish and fish larvae on passage through some regulating structures like undershot weirs.

Fish community health can be improved through targeted management actions and re-instatement of appropriate water regimes. Some complementary measures to assist fish populations have recently occurred with the construction of a thermal (cold water) curtain on Burrendong Dam and the construction of fishways at a number of weirs and regulators (NSW DPI 2015).

Managing flows to improve native fish populations involves restoring a range of flows and providing access to physical (wetted) habitat to allow movement and improve the carrying capacity of waterways and wetlands. Fish population structure is closely tied to the frequency of successful breeding events and enhanced recruitment, as well as improved maintenance, condition and movement outcomes for all life-history stages.

Objectives and targets for native fish in the Macquarie-Castlereagh WRP relate to increasing distribution and abundance of selected species and ensuring a stable population structure that includes representation of young-of-year, juvenile and adult-life-history stages (Table 1). These objectives can be achieved by providing flows across the entire spectrum of the flow regime (from low flows through to overbank flow events) that meet the EWRs for all fish species.

Table 1 Native fish ecological objectives

LTWP Objective		Target fish species	LTWP Targets ¹		
			5 years (2024)	10 years (2029)	20 years (2039)
NF1	No loss of native fish species	All native species	All known species detected annually		
NF2	Increase the distribution & abundance of short to moderate-lived generalist native fish species	Australian smelt, carp gudgeon, flat-headed gudgeon, bony herring, Murray-Darling rainbowfish, unspotted hardyhead, dwarf flat-headed gudgeon, mountain galaxias	Increased distribution & abundance of short to moderate-lived species compared to 2014 assessment No more than 1 year without detection of immature fish (short-lived) No more than 2 years without detection of immature fish (moderate-lived species)		
NF3	Increase the distribution & abundance of short to moderate-lived floodplain	Olive perchlet ³ , flathead galaxias			

¹ For annual detection targets, annual detection is at a fish monitoring zone scale, which may not correspond to PUs. Unless otherwise stated the targets are for detection in relevant PUs/ NSW Fisheries Basin Plan Environmental Outcomes Monitoring Zones where fish are expected to occur.

³ In the Macquarie catchment, the olive perchlet is considered to best fit the floodplain-specialist functional group and the purple-spotted gudgeon, the riverine specialist functional group, however both have some characteristics of both functional groups.

LTWP Objective		Target fish species	LTWP Targets ¹		
			5 years (2024)	10 years (2029)	20 years (2039)
	specialist native fish species ²				
NF4	Improve population structure for moderate to long-lived flow pulse specialist native fish species	Golden perch, silver perch, spangled perch, Hyrtl's tandan	Juvenile & adult fish detected annually No more than 2 consecutive years without recruitment in moderate-lived species No more than 4 consecutive years without recruitment in long-lived species		
NF5	Improve population structure for moderate to long-lived riverine specialist native fish species	Murray cod, northern river blackfish, eel-tailed catfish, purple-spotted gudgeon			
			Minimum of 1 significant recruitment event in 5 years ⁴	Minimum of 2 significant recruitment events in 10 years ⁴	Minimum of 4 significant recruitment events in 20 years ⁴
NF6	A 25% increase in abundance of mature (harvestable sized) golden perch & Murray cod	Golden perch, Murray cod	Length-frequency distributions include size classes of legal take size for golden perch & Murray cod 25% increase in abundance of mature golden perch & Murray cod		
NF7	Increase the prevalence &/or expand the population of key short to moderate-lived floodplain specialist native fish species into new areas (within historical range)	Flathead galaxias, olive perchlet ²	Adults detected annually in specified PUs No more than 1 year (short-lived), 2 years (moderate-lived species) or 4 years (long-lived species) without detection of immature fish in specified PUs		
NF8	Increase the prevalence &/or expand the population of key moderate to long-lived riverine specialist native fish species into new areas (within historical range)	Trout cod, eel-tailed catfish, purple-spotted gudgeon, river blackfish, Macquarie perch			
				Increased distribution & abundance in specified PUs	

² In many cases conservation stocking will be required to achieve range expansion to historical areas for floodplain specialist species.

⁴ Young-of-year comprise more than 30% of the population.

LTWP Objective		Target fish species	LTWP Targets ¹		
			5 years (2024)	10 years (2029)	20 years (2039)
NF9	Increase the prevalence and/or expand the population of key moderate to long-lived flow pulse specialists native fish species into new areas (within historical range)	Hyrti's tandan			Increased distribution & abundance in specified planning units



Figure 4 Murray cod. Photo Gunther Schmida

3.2 Native vegetation objectives

The riverine, wetland and floodplain environments of the Macquarie-Castlereagh support a diverse range of water-dependent vegetation communities. Native water-dependent vegetation provides important food and habitat for many animals and plays an important role in maintaining water quality, stabilising river banks and contributing organic matter to feed the ecosystems of rivers and wetlands.

This Plan focuses on the riverine, wetland and floodplain vegetation communities that depend on permanent and periodic inundation from the river. These are primarily river red gum, black box and coolibah forests and woodlands, lignum shrublands and non-woody wetland vegetation communities—such as reedbeds and water couch meadows—sometimes referred to as ‘semi-permanent wetland vegetation’.

River red gum forest and woodlands

River red gum forests and woodlands are an important component of the Macquarie-Castlereagh region's rivers and wetlands, supporting ecological functions and providing important habitat. River red gum forest and woodlands have a life-cycle that is heavily dependent upon regular, periodic, but not permanent inundation.

The Macquarie Marshes supports the largest river red gum area in the northern Murray-Darling Basin (approximately 40,000 hectares). River red gum communities have declined in condition and extent in the Macquarie Marshes since 1991 (Bowen & Simpson 2010; DECCW 2010).

To maintain and improve the condition of river red gum forests and woodlands in the Macquarie Marshes, efforts need to be made to inundate areas in low and moderate condition to ensure they remain in the landscape (Bowen and Simpson 2010).

Coolibah and black box woodlands

Coolibah and black box woodland communities are dependent on periodic inundation and are generally located on higher parts of the floodplain throughout the northern and western parts of the Macquarie-Castlereagh. Both communities are part of the coolibah-black box endangered ecological community listed under the *NSW Biodiversity Conservation Act 2016*. The extent and condition of coolibah-black box communities has declined since 1991 due to clearing and altered flow regimes (Bowen & Simpson 2010).

Lignum shrublands

Lignum shrubland is a relatively limited community present in some northern and western parts of the WRPA. This multi-stemmed shrub grows primarily in floodplain and wetland areas and provides important habitat for a range of species including colonially-nesting waterbirds. Lignum shrubland has reduced significantly in area, with a reduction of 89% recorded from 1991–2008 (Bowen & Simpson 2010).

Non-woody wetland vegetation

The Macquarie Marshes contain significant stands of common reed (4,800 hectares), cumbungi (1,500 hectares), water couch (5,200 hectares) and mixed sedgelands (6,500 hectares) that require regular, frequent and prolonged inundation. These communities provide important foraging and breeding habitat for waterbirds. Their extent of these communities vary according to prevailing inundation conditions.

Objectives

The focus of the native vegetation objectives in this Plan are to maintain extent of flow-dependent vegetation communities. In some areas of the *actively managed floodplain* it may be possible to improve the condition of these communities over time, and objectives for these have been set accordingly (Table 2).

Table 2 Native vegetation ecological objectives

LTWP Objective			Targets		
			5 years (2024)	10 years (2029)	20 years (2039)
NV1	Maintain or improve the extent & viability of non-woody vegetation communities occurring within channels ⁵		Increase cover of non-woody, inundation-dependent vegetation within or closely fringing river channels following inundation events (evaluated at specific sites only)		
NV2	Maintain or increase the extent & viability of non-woody vegetation communities in wetlands & on floodplains		<p>Maintain the extent of non-woody, inundation dependent vegetation to 2017 OEH amalgamated mapped extent⁶</p> <p>Successful growth, flower & seed set of common reed, cumbungi & water couch at least 2 years in any 5-year period at key wetland sites to ensure long term persistence & viability (evaluated at specific sites only)</p>		
NV3	Maintain the extent & improve the condition of river red gum communities closely fringing river channels ⁷		<p>Maintain the extent of river red gum communities closely fringing river channels</p> <p>Over a 5-year rolling period, maintain the proportion of river red gum communities closely fringing river channels that are in moderate or good condition⁸</p> <p>Over a 5-year rolling period, no further decline in the condition of river red gum communities closely fringing river channels that are in poor or degraded condition⁸</p>		
NV4a	Maintain or increase the extent & maintain or improve the condition of native forest, woodland & shrubland communities on floodplains ⁷	River red gum forest	Maintain the 2017 OEH amalgamated mapped extent of river red gum forest & woodland & black box & coolibah woodland communities	Maintain the 2017 OEH amalgamated mapped extent of river red gum forest & woodland & black box & coolibah woodland communities	
NV4b		River red gum woodland	Over a 5-year rolling period, maintain the extent & proportion of river red gum forests & woodlands & black box & coolibah woodlands in moderate or good condition ⁸	Over a 5-year rolling period, maintain or increase the proportion of river red gum forests & woodlands & black box & coolibah woodlands in moderate or good condition ⁸	
NV4c		Black box woodland			

⁵ In the Macquarie there is no significant non-woody vegetation outside the channel which is not in wetlands (covered by NV2). This applies to in-channel vegetation in the Macquarie.

⁶ Measured over a 5-year rolling period to account for variation between naturally dry and wet times.

⁷ Held environmental water cannot generally be delivered at the high flow levels required to meet some of the key needs of riparian river red gum vegetation communities (e.g. ground-cover condition), for woodlands and shrublands outside the actively managed floodplain, nor for some channel forming processes. Planned environmental water services these needs. Policy changes which affect these flows will risk the health of these vegetation communities and processes.

⁸ Measured by MDBA Stand Condition tool.

LTWP Objective		Targets		
		5 years (2024)	10 years (2029)	20 years (2039)
NV4d	Coolibah woodland	Over a 5-year rolling period, no further decline in the condition of river red gum forests & woodlands & black box & coolibah woodlands in poor or degraded condition ⁸ Maintain or improve the age class structure of river red gum, black box & coolibah communities (measured in selected sites only)	Over a 5-year rolling period, no further decline in the condition of river red gum forests & woodlands & black box & coolibah woodlands in poor or degraded condition ⁸ Maintain or improve the age class structure of river red gum, black box & coolibah communities (measured in selected sites only)	
NV4e	Lignum shrublands	Maintain the 2017 OEH amalgamated mapped extent of lignum shrubland communities Over a 5-year rolling period, increase the proportion of lignum communities in intermediate to good condition, on the <i>actively managed floodplain</i> , & maintain the proportion on the remaining floodplain Over a 5-year rolling period, no further decline in the condition of lignum shrublands in poor condition		

3.3 Waterbird objectives

Waterbirds respond to variable climatic conditions and are known to travel great distances in search of ideal wetland habitat. The total number and diversity of waterbird species fluctuates with seasonal conditions, with widespread flooding generated by high rainfall in the upper catchment an important trigger for large colonial waterbird breeding events. Aspects of flows that are considered important to the ecological requirements of waterbirds include, timing, inundation duration and extent, the rate of rise and fall of water, depth of inundation, frequency of events and inter-event dry periods (Ralph & Rogers 2011).

The Macquarie-Castlereagh has records from 1992–2012 for 11 waterbird species (not including vagrants) listed under one or more international migratory bird agreements (JAMBA, CAMBA and RoKAMBA). These are the Caspian tern, black-tailed godwit, common greenshank, common sandpiper, curlew sandpiper, Latham's snipe, marsh sandpiper, red-necked stint, sanderling, sharp-tailed sandpiper and wood sandpiper (OEH 2012). There are also records for the Macquarie-Castlereagh from 1992–2012 for 11 waterbird species listed under the NSW Biodiversity Conservation Act, including Australasian bittern, blue-billed duck, freckled duck, magpie goose and brolga. Most of these species are dependent upon wetlands and floodplains.

The Macquarie Marshes provide important habitat for a rich diversity of waterbird species. Seventy-six waterbird species (44 breeding) have been recorded in the Marshes (Kingsford & Auld 2005). They regularly support more than 20,000 waterbirds, and in times of large floods, numbers can exceed more than 500,000 waterbirds.

The Macquarie Marshes are highly significant for colonial nesting waterbirds, particularly the number of nests, frequency of breeding and the diversity of species. Colonies are generally reliant upon dam spill events that occur during winter, spring or summer. During these events, the Marshes can support colonies totalling more than 100,000 waterbird nests (Spencer 2017). Recent examples of large-scale events include 2000, 2010/11 and 2016/17, with smaller breeding events in 2011/12 (OEH 2012, Spencer et al. 2018). The Marshes are especially significant for supporting large breeding colonies of straw-necked ibis, intermediate egret, rufous night-heron and royal spoonbill (Spencer et al. 2018) and for being one of only a few sites in NSW where the magpie goose breeds (OEH 2012).

The Macquarie Marshes also act as a refuge in dry periods by providing habitat for waterbirds when many other inland wetlands and waterways have dried out (OEH 2012).

In the 30 years to 2012, annual waterbird surveys revealed a 72% decline in average waterbird abundance across the Murray-Darling Basin (MDBA 2014a). This is a critical observation because waterbirds are an important indicator of wetland health as their abundance and diversity are related to the total area of wetland available, the health of wetland vegetation and the abundance of food resources e.g. fish and aquatic vegetation (Kingsford 1999).

Prior to regulation of the Macquarie River in 1967, colonial waterbirds bred regularly in the Macquarie Marshes; however, river regulation has reduced the frequency of larger flow events reducing opportunities to breed. River regulation and clearing of native vegetation has impacted colonial waterbird breeding habitats, with many breeding locations in the Marshes considered to be in poor condition (OEH 2010). These factors have led to significant declines in diversity, abundance and number of breeding events and breeding sites used (OEH 2013).

The Macquarie Marshes has been identified as a key wetland where restoration of habitat for waterbirds, and reinstatement of flow regimes that mimic natural, can contribute to recovery of waterbird populations both in-catchment and across the Murray-Darling Basin (MDBA 2014a).

Objectives

Objectives and targets in this plan for waterbirds relate to maintaining diversity and supporting waterbird population recovery compared to the 2012–2016, or, where stated 1992–2012, baseline (Table 3). This will be sought mainly through maintaining and, where possible, improving key waterbird breeding and foraging habitat, and through the delivery of targeted flows to maintain or augment naturally initiated colonial nesting events through to post-fledging care, such as the 2016/17 summer colonial nesting event.



Figure 5 Intermediate egret in breeding plumage. Photo John Spencer OEH

Table 3 Waterbird ecological objectives

LTWP Objective ⁹		Targets		
		5 years (2024)	10 years (2029)	20 years ¹⁰ (2039)
WB1	Maintain the number and type of waterbird species ¹¹	Maintain a 5-year rolling average of 41 or more waterbird species across the 5 functional groups in the Macquarie Marshes		
			Identify at least 62 waterbird species in the Macquarie Marshes in a 10-year period	Identify at least 69 waterbird species in the Macquarie Marshes in a 20-year period
WB2	Increase total waterbird abundance across all functional groups	Total abundance of the 5 functional groups maintained in the Macquarie Marshes compared to the 5-year 2012–16 period	Total waterbird abundance increased by 20–25% in the Macquarie Marshes compared to the 5-year 2012–16 period, with increases in all functional groups	Maintain or increase total waterbird abundance in the Macquarie Marshes compared to the 10-year target, with increases in all functional groups
WB3	Increase opportunities for non-colonial waterbird breeding	Total abundance of non-colonial waterbirds in the Macquarie Marshes maintained compared to the 5-year 2012–16 baseline & breeding recorded in at least 14 non-colonial waterbird species over the 5-year period	Total abundance of non-colonial waterbirds in the Macquarie Marshes increased by 20–25% compared to the 5-year 2012–16 baseline with breeding detected in at least 14 non-colonial waterbird species over the 5-year period	Maintain or increase total abundance of non-colonial waterbirds in the Macquarie Marshes compared to the 10-year target, with breeding detected in at least 14 non-colonial waterbird species in any 5-year period
WB4	Increase opportunities for colonial waterbird breeding	Support active waterbird colonies in the Macquarie Marshes by maintaining the water depth & duration of inundation (as required) to support breeding through to completion (from egg laying through to fledging including post-fledgling care) & maintain duration of inundation in key foraging habitats to enhance breeding success & the survival of young		
WB5	Maintain the extent & improve condition of waterbird habitats	Maintain extent & improve condition of nesting vegetation, including common reed, lignum, cumbungi, river red gum & river cooba, in known colonial breeding locations in the Macquarie Marshes		
		Maintain or increase extent & improve condition of waterbird foraging & breeding locations in the Macquarie Marshes (to be evaluated under targets set for native vegetation)		

⁹ Because most waterbird surveys in the Macquarie-Castlereagh have occurred in the Macquarie Marshes, information on waterbird diversity and abundance is more reliable for these areas than throughout the rest of the catchment. The objectives for waterbirds described in this LTWP are therefore only provided for the wetland areas where progress toward meeting objectives can be evaluated and contribute the most towards the Basin - scale outcomes of the Basin Plan.

¹⁰ Targets will be further refined from 2029 onwards following additional data collection.

¹¹ Not including vagrant species, targets based on 1992–2012 baseline.

3.4 Priority ecosystem function objectives

A number of key functions and processes have been identified in the Macquarie Castlereagh and these are described below.

Drought refuge for water-dependent species

Instream pools and floodplain lagoons are extremely valuable refuges in riverine landscapes. Other types of instream refugia include logs, wet undercut banks, riffles, sub-surface stream sediments and riparian vegetation (Boulton 2003). Refugia are critical to the survival of many aquatic species during dry spells and drought, and act as source populations for subsequent recolonisation and population growth (Adams & Warren 2005; Arthington et al. 2005).

Quality instream habitat

The physical form of instream habitats, including the location of riparian and instream vegetation, channel shape and bed sediment, is influenced by river flow. For example, fresh and bankfull flows with sufficient velocity are required to maintain pool depth and riffles by scouring out bed material and initiating material transportation downstream.

Movement and dispersal opportunities for aquatic biota

Longitudinal and lateral connectivity allows organisms to move and disperse between environments. It can be essential for maintaining population viability (Amtstaetter et al. 2016) by allowing individuals to move to different habitat types for breeding and conditioning, and recolonisation following disturbances like flood and drought. Flow pulses can promote dispersal of early life stages for a range of species from the breeding site and promote genetic diversity among catchments (Humphries & King 2004). Movement is currently limited in many reaches by in-stream structures.

Instream and floodplain productivity

The supply of organic material underpins all river food webs by providing the food energy needed to drive life. The sources of organic material, the timing of its delivery and how long it remains in a section of river depend very much on the flow regime and the nature of the riparian vegetation.

River flow management can be used to increase carbon and nutrient sources in-channel by increasing the frequency of floodplain inundation. Re-wetting patches (e.g. river channels, channel benches, floodplains following drying provides a pulse of terrestrial carbon available for potential use by consumers (e.g. Lanhans & Tockner 2006) and the flow of water enhances the physical breakdown of leaves, branches and other terrestrial detritus (Mora-Gomez et al. 2015).

Groundwater-dependent biota

While this Plan is primarily focused on the management of surface water, the Macquarie-Castlereagh Groundwater Source plays an important ecological role in supporting terrestrial and aquatic ecosystems, particularly during extended dry periods where they can be critical for maintaining refuges.

Groundwater is predominately found in alluvial sediments on the plains in the lower catchment, with the highest yielding aquifers located north-west of Narromine. The Great Artesian Basin underlies the northern part of the Macquarie catchment downstream of Warren (MDBA, n.d.).

To continue to support groundwater dependent ecosystems in the Macquarie-Castlereagh, objectives relate to maintaining the mapped extent of groundwater-dependent vegetation communities and groundwater levels within the natural range of variability over the long-term.

Sediment, carbon and nutrient exchange

The frequency of flows that connect rivers with their riparian corridors and floodplains has been substantially reduced. Water volumes released from Burrendong Dam typically do not exceed channel capacity. The loss of lateral connectivity between rivers and their floodplains has altered water movement, the flux of sediment, nutrients, carbon, and biota from and to the river (Baldwin et al. 2016). Consequently, the amount of dissolved organic carbon entering the main channels is reduced because of less frequent wetting of benches, flood runners and floodplains (Westhorpe et al. 2010). Longitudinal connectivity is equally important and fulfils the important environmental function of transporting nutrients and sediments between environments (MDBA 2014a).

Inter-catchment flow contributions

Longitudinal connectivity between catchments fulfils important environmental functions e.g. moving nutrients and sediments, allowing for organisms to disperse and improving water quality. River regulation and diversions have reduced the volume of water passing into downstream catchments.

Connectivity between key PUs and provision of end of system flows will contribute to improved environmental outcomes in the Macquarie-Castlereagh and Barwon-Darling catchments.

Table 4 Ecosystem function objectives

Objective		Description & key contributing processes	Targets		
			5 years (2024)	10 years (2029)	20 years (2039)
EF1	Provide & protect a diversity of refugia across the landscape	Water depth & water quality in pools (in-channel), core wetlands & lakes Condition of vegetation in core wetlands & riparian zones	Core wetland habitats ¹² are protected ¹³ during dry times Cease-to-flow periods do not exceed maximum durations as specified in PU EWRs Adequate water depth is maintained in key refuge pools ¹⁴ during dry times In key refuge pools ¹⁴ maintain daily average dissolved oxygen >4 mg/L & hourly levels >2 mg/L		
EF2	Create quality instream, floodplain & wetland habitat	<ul style="list-style-type: none"> Regulation of dissolved oxygen, salinity & water temperature Flow variability & hydrodynamic diversity Provision of diverse wetted areas Appropriate wetting & drying cycles Geomorphic (erosion/deposition) processes that create & maintain diverse physical habitats Appropriate rates of fall to avoid excessive bank erosion Control of woody-vegetation encroachment into river channels & wetlands 	Recession management: Rate of fall does not exceed the 95th percentile of natural rates ¹⁵ during regulated water deliveries Minimum flow variability: Period for which small & large freshes are held at constant level ($\pm 5\%$) does not exceed 20 days to avoid bank slumping & support in-stream function ¹⁶ Channel form: Watering requirements for overbank flows are met (refer to fish/veg/bird watering requirements) ⁷ Bench & pool formation & fine sediment scouring: Watering requirements for freshes are met (refer to fish/veg/bird watering requirements) Create hydrodynamic complexity for large-bodied fish: Flows with velocities of 0.3 to 0.4 m/s provided as per watering requirements for freshes (refer to fish watering requirements)		

¹² Core wetland habitats are those identified by Thomas et al. (in prep) as the 'purple zone' in the Macquarie Marshes, which is inundated with an average return interval of one to two years.

¹³ Maintained in a state that will provide habitat for biota and allow recovery to good condition with a wet season.

¹⁴ To be identified – see Table 19. Monitored at selected sites only.

¹⁵ Calculation of these percentiles will be required for relevant planning units. 'Natural' rates estimated from pre-1966 observed data where a multi-decadal record exists or, where this is not available, modelled natural data.

¹⁶ Note: There may be some instances where, for ecological reasons, relatively constant flows are required, such as keeping water levels below nesting colonies. In these cases exceptions may be required at the relevant planning unit. However, even these deliveries may still be able to be subtly altered to provide variability.

Objective		Description & key contributing processes	Targets		
			5 years (2024)	10 years (2029)	20 years (2039)
EF3	Provide movement & dispersal opportunities within catchments for water-dependent biota to complete lifecycles: a. within catchment b. between catchments	<ul style="list-style-type: none"> Dispersal of eggs, larvae, propagules & seeds downstream & into off-channel habitats Migration to fulfil life-history requirements Foraging of aquatic species Recolonisation following disturbance 	Increase, compared to 2004–2017 ¹⁷ , dispersal opportunities between sub-catchments & between river reaches for moderate to long-lived flow riverine & pulse specialist native fish through key fish passages ¹⁸ .		
			Long-term observed frequency of connectivity sufficient to allow native fish movement, with a focus on moderate to long-lived flow pulse specialists, between the Barwon River & the Lower Macquarie & Lower Marthaguy is improved, & between the Barwon River & the unregulated Castlereagh & Bogan Rivers & Marra Creek are maintained ¹⁶ , compared to the 2004–2017 baseline scenario		
EF4	Support instream & floodplain productivity	<ul style="list-style-type: none"> Aquatic primary productivity (algae, macrophytes, biofilms, phytoplankton) Terrestrial primary productivity (vegetation) Aquatic secondary productivity (zooplankton, macroinvertebrates, fish larvae, adult fish) Decomposition of organic matter 	Maintain or increase the proportion of wetland & floodplain vegetation that is in good condition ¹⁹ over a 5-year rolling period		
			No decline in key native fish species ²⁰ condition metrics	Improve key native fish species condition metrics	
			Maintain the abundance & distribution of decapod crustaceans	Improve the abundance & distribution of decapod crustaceans	

¹⁷ To be assessed against a combination of observed data and the modelled baseline scenario which represents the consumptive use and the rules and sharing arrangements as at June 2009. Once environmental flow behaviours are updated in the models, this version should be used. Comparisons will need to take into account any limitations in the model and, for the observed data, the comparability of the weather during the baseline period and target period.

¹⁸ Key fish passages to be identified by NSW Fisheries.

¹⁹ In line with condition targets set for the native vegetation objectives.

²⁰ Key fish species that are relevant in each planning unit, as described in the targets for the native fish objectives.

Objective		Description & key contributing processes	Targets		
			5 years (2024)	10 years (2029)	20 years (2039)
EF5	Support nutrient, carbon & sediment transport along channels, & between channels & floodplains/wetlands	<ul style="list-style-type: none"> Sediment delivery to downstream reaches & to/from anabranches, floodplains & wetlands Mobilisation of carbon & nutrients from in-channel surfaces (e.g. benches/banks), floodplains & wetlands & transport to downstream reaches & off-channel habitats Dilution of carbon & nutrients that have returned to rivers 	Maintain the frequency & duration of events that drive nutrient & carbon (DOC) processes (at selected evaluation sites) along channels (freshes & overbank events) Maintain extent & condition of floodplain vegetation (measured under vegetation theme targets)		
EF6	Support groundwater conditions to sustain groundwater-dependent biota	<ul style="list-style-type: none"> Groundwater recharge & discharge Dilution of saline/acidic groundwater Salt export from the Murray-Darling Basin 	Maintain the current mapped extent of groundwater-dependent vegetation communities & specifically maintain the freshwater lens in Macquarie Marshes ²¹ Maintain groundwater levels within the natural range of variability over the long-term (monitored in selected sites only)		
EF7	Increase the contribution of flows into the Murray & Barwon-Darling from tributaries	<ul style="list-style-type: none"> Provision of end of system flows to support ecological objectives in downstream catchments 	End of system flow volumes to the Barwon River increase by 10% compared to 2004–2017 ¹⁷		

²¹ Note: Groundwater systems in the Marshes (and generally) are not well understood and there may be a need for more information gathering before this can be meaningfully measured. These targets are one way of measuring but may not be able to be done in the first instance. In the meantime, it is known that large floods recharge the groundwater systems and flush salts from the soils. These are not able to be delivered with held environmental water. Hence there is a need to protect these larger events when they do occur.



Figure 6 Salmon striped frog. Photo: Joanne Ocock/OEH

3.5 Objectives for other species (flow-dependent frogs)

The delivery of environmental watering requirements for waterbirds, native fish, vegetation and functions/processes will benefit other species. However, there are some flow-dependent species that are significant (due to their vulnerability, cultural significance or importance in the food chain) and which have watering requirements that are different to those covered by other themes.

In the Macquarie this is relevant for frogs. Frogs are an important food source for waterbirds, fish and reptiles. Flow-dependent frogs have similar watering requirements to waterbirds for breeding, however, they are not highly mobile, so they require refuge between watering events. As such, the inclusion of flow-dependent frog objectives places greater emphasis on the need to maintain refuge (including permanent waterbodies for some species) in the floodplain.

Nearly half of the frog species found in the floodplain wetlands of the Murray-Darling Basin are responsive to flows and have been categorised as 'flow-dependent' (OEH, unpublished). There are six flow-dependent frog species found in the Macquarie Marshes, and their populations depend on wetland inundation of sufficient extent, duration and seasonality (Ocock and Spencer 2018), and on the condition of wetland habitat for feeding, breeding and refuge.

Table 5 Objectives for other species (flow-dependent frogs)

Ecological objectives		Targets ²²		
		5 years (2024)	10 years (2029)	20 years (2039)
OS1	Maintain species richness & distribution of flow-dependent frog communities	Over a 5-year rolling period, detect, in each assessment period, all 6 flow-dependent frog species known from the Macquarie Marshes based on comprehensive surveys over the period 2015–2017		
OS2	Maintain successful ²³ breeding opportunities for flow-dependent frog species	Over a 5-year rolling period, maintain proportion of wetlands sites where breeding activity ²⁴ of flow-dependent frog species is detected in the Macquarie Marshes compared to the 2015-17 period		

²² Baseline data from Ocock and Spencer (2018).

²³ Successful relates to opportunities for species to complete breeding life cycle, i.e. laying eggs, to development of tadpoles through to metamorphs (juvenile frogs), which relates to water requirements for minimum duration of inundation.

²⁴ We define breeding activity as observations of male frog callings, frog spawn, tadpoles and/or recently metamorphosed juvenile frogs. These observations are evidence of potential recruitment of new individuals into the breeding population.

4. Environmental water requirements

A river's flow or inundation regime influences the ecological characteristics of that river's ecosystems (Poff and Zimmermann 2010). A flow regime represents the totality of flow events. Individual flow events shape river channels, provide cues for key biological processes such as breeding or migration, support dispersal of plants and animals and shape how a river links with its floodplain. However, it is the overall flow that dictates whether populations are stable, provides ecosystem resilience and ultimately our long-term outcomes.

Flow components can be defined ecologically or hydrologically and can be partitioned into flow types or flow components—such as base flows, freshes, overbank flows—that describe the height or level of the flow within a river channel or its extent across a floodplain (see Figure 7).

Each flow component can provide for a range of ecological roles. For example, a small fresh might inundate river benches that provide access to food for native fish and support in-channel vegetation. Similarly, an overbank flow may support carbon exchange between the river and its floodplain and improve river red gum condition. Flow rate bands for flow components in the Macquarie-Castlereagh are shown in **Error! Reference source not found.**

An environmental watering requirement (EWR) is the flow or inundation regime that a species, population or community needs to ensure its survival and persistence. EWRs are based on current scientific information of species' biological and ecological needs, such as what it needs to feed, breed, migrate and disperse.

Meeting the lifetime needs of an aquatic organism (plant or animal) might require a combination of several different flow components over time. For example, a native fish species may require a 'small fresh' as a 10-day pulse in late winter to cue spawning, followed by a relatively stable flow for 2–4 weeks in early spring to support nesting. Once the fish reaches maturity (1–3 years) it may require a fast-flowing river in combination with 'overbank' flows to trigger dispersal and migration.

4.1 Developing EWRs

Development of EWRs for LTWPs was an iterative process that started with an assessment of watering requirements of individual species, then of guilds or functional groups. Where EWRs overlapped between species or groups, the EWRs were combined to provide a single EWR for a flow component.

At the PU-scale, EWRs were informed by an understanding of the channel morphology and hydrology. This included an analysis of channel cross-sections, floodplain inundation data, observed flow data, modelled flow data and operational experience.

Important flow regime characteristics to meet life-history needs and each of the LTWP objectives are described in Table 9. The combined EWRs, grouped by flow component, for all biota and functions in the Macquarie-Castlereagh WRP are presented in Table 8. Each EWR is expressed as a flow component that has been assigned an ideal timing, duration and frequency based on the suite of plants, animals and functions it supports. They identify ideal minimum durations and frequencies and maximum periods between events.

A detailed description of the methods (in prep.) used to develop LTWPs can be found at www.environment.nsw.gov.au.

The health of the aquatic and floodplain ecosystem is dependent on the full flow regime across every day of every season of every year. All flow types (baseflows, freshes and overbanks), their timings, durations, frequencies and the gaps between them dictate the

environment which is available to the plants and animals which depend on the river. In Table 8 there are a number of key flow requirements (EWRs). However, these are only select parts of the flow regime. The river's flow regime is greater than the sum of these parts. Any evaluation of the potential effect of changes to the river should not only assess the EWRs, but also other measures of flow, including measurements of variability, seasonality and overall flow volumes (particularly to the end of system and the Macquarie Marshes). Additionally, for the EWRs, we have identified ideal minimum durations and frequencies and maximum periods between events. Achieving these does not signal that no problems may exist. Where a change causes reductions in EWRs, even if the minimums are still achieved, this may still be an important decline in the flow regime.

OEH acknowledge that achieving EWRs alone does not mean objectives will be reached. There are a range of complimentary measures that will improve chances of achieving objectives, and other variables such as climate change, weather and land use will impact upon the environmental outcomes sought.




Figure 7 A simplified conceptual model of the role of flow regime components

Table 6 Description of function role provided by flow components

Flow component	Description
Overbank/ wetland inundating flow	<p>Both overbank & wetland inundation flows provide broad scale lateral connectivity with floodplain & wetlands. Supports nutrient, carbon & sediment cycling between floodplain & channel. Promotes large-scale productivity.</p> <p>Overbank flows are used to describe flows when they are above bankfull.</p> <p>Wetland inundation flows are used to describe:</p> <ul style="list-style-type: none"> flows that fill wetlands via regulating structures below bankfull over weeks or sometimes months (i.e. longer than a typical fresh/pulse), or flows that are required to inundate wetlands in areas where there are very shallow channels or no discernible channels exist (e.g. terminal wetlands).
Bankfull flow	<p>Inundates all in-channel habitats & connects many low-lying wetlands. Partial or full longitudinal connectivity. Drown out of most small in-channel barriers (e.g. small weirs).</p>

Flow component	Description
Large fresh (pulse)	Inundates benches, snags & inundation-tolerant vegetation higher in the channel. Supports productivity & transfer of nutrients, carbon & sediment. Provides fast-flowing habitat. May connect wetlands & anabranches with low commence-to-flow thresholds.
Small fresh (pulse)	Improves longitudinal connectivity. Inundates lower benches, bars, snags & in-channel vegetation. Trigger for aquatic animal movement & breeding. Flushes pools. May stimulate productivity/food webs.
Baseflow	Provides connectivity between pools & riffles & along channels. Provides sufficient depth for fish movement along reaches.
Very low flow	Minimum flow in a channel that prevents a cease-to-flow. Provides connectivity between some pools & prevents stratification of pools.
Cease-to-flow	Partial or total drying of the channel. Stream contracts to a series of disconnected pools. No surface flow.

4.2 Flow threshold estimates for flow components in the regulated Macquarie-Castlereagh

Table 7 Flow threshold estimates for flow components in the regulated Macquarie-Castlereagh WRP

Note: All units ML/d, except total volume which is shown as ML.

Planning unit	Gauge	Low flows		Fishes		Overbank		
		Very low	Baseflow	Small	Large	Small	Medium	Large
REGULATED CUDGEGONG RIVER								
Cudgegong River @ Yamble Bridge	421019	40	40	200	1,000	12,000 (not deliverable) ⁷	Not applicable – PU not characterised by extensive river red gum forests on the wider floodplain	
REGULATED MACQUARIE RIVER								
Burrendong to Baroona (@Dubbo)	421001	200	200	500	6,000	65,000 (not deliverable) ⁷	Not determined – non-riparian river red gum forests & woodlands are less extensive in this PU	Not applicable – PU not characterised by extensive black box or coolibah woodlands on the wider floodplain
Baroona to Warren (@ Warren Weir)	421004	200	200	450	4,000	12,000 (not deliverable) ⁷	16,000	18,000
Warren to Macquarie Marshes (@combined Marebone Break & Macquarie River below Marebone gauges)	421088 + 421090	30	100	350	2,500	4,000	5,500	Not determined

Planning unit	Gauge	Low flows		Freshes		Overbank		
		Very low	Baseflow	Small	Large	Small	Medium	Large
MACQUARIE MARSHES								
Northern and Southern river flow thresholds (Macquarie River downstream Marebone & Oxley Station)	421090, 421022	25	65	300	1,000	2,900@ downstream Marebone 2600 @ Oxley Station		
Whole of Macquarie Marshes wetland flows – Northern, Southern & Eastern Marshes (@combined Macquarie River & Marebone Break below Marebone gauges)	421088 + 421090					For core wetlands (small overbank/ wetland 1): 60,000 ML over 90 days For wetland non-woody vegetation (small overbank/ wetland 4): 100,000 ML within 90 days	250,000 ML within 120 days	440,000 ML within 150 days
Eastern Marsh wetland flow (Gum Cowal @ Bifurcation)	421146						8,000 ML on days above 400 ML/d, within 40-day period	Covered by whole of Marshes flow – see above.
LOWER MACQUARIE RIVER								
Lower Macquarie (@ Bells Bridge (Carinda))	421012	10	65	140	700	1,900	Not determined	Not determined
LOWER MARTHAGUY CREEK								
Marthaguy Creek (@ Carinda)	421011	20	20	70	800	2,900	Not determined	Not determined




Figure 8 Schematic of the main watercourses and streamflow gauges used for EWRs (Not to scale)

4.3 Catchment-scale EWRs for Macquarie-Castlereagh

Table 8 Catchment-scale EWRs for Macquarie-Castlereagh WRPA

Flow component		Environmental objectives	Timing ²⁵	Duration ²⁶	Frequency ²⁷	Maximum inter-event period ²⁸	Other requirements
Cease-to-flow	CF1	Native Fish: NF1 – Survival & condition (all species) Ecosystem Functions: EF1, 2 – refuge habitat	In line with historical low season flow ²⁹	In line with natural, unless key refuges endangered	No greater than natural ²⁹	N/A	When restarting flows, avoid harmful water-quality impacts, such as de-oxygenated refuge pools.
Very-low flow	VF1	Native Fish: NF1 - Survival & condition (all species) Ecosystem Functions: EF1, 2 – refuge habitat	Any time	No less than natural ²⁹	No less than natural ²⁹	No greater than natural ²⁹	Flow ideally >0.03–0.05 m/s to de-stratify refuge pools
Baseflow	BF1	Native Fish: NF1, 2, 3, 4, 5, 6, 7, 8, 9 – condition & movement (all species) Native Vegetation: NV1 – in-channel Ecosystem Functions: EF1, 2, 3a Other species (frogs): OS1	Any time	No less than natural ²⁹ – maintain refuge habitat & support connectivity	No less than natural ²⁹	As required during dry periods	Minimum depth of 0.3 m to allow fish passage

²⁵ Recommended ideal timing for native fish is linked to maximising spawning and recruitment outcomes based on known spawning seasons. Ideal timings where the purpose is for dispersal and/or productivity have also been recommended; however, these may occur anytime with a movement and/or condition outcome still expected from native fish.

²⁶ Recommended minimum duration for native fish is linked to maximising spawning and recruitment outcomes based on known egg hatch time and morphology.

²⁷ Recommended ideal frequency for native fish is linked to providing conditions that protect and improve native fish populations in heavily impacted systems. To achieve this recovery more frequent events that maximise native fish outcomes may be required.

²⁸ Recommended ideal period between events for spawning and recruitment objectives is linked to the longevity of species, providing a guide to the maximum period between these outcomes before risk of significant population decline. Objectives related to in-channel dispersal and/or productivity have been set at annual frequencies to provide conditions that protect and improve native fish populations in heavily impacted systems.

²⁹ Determined using pre-1966 observed data where a multi-decadal record exists or modelled natural where sufficient observed data lacking.

Flow component		Environmental objectives	Timing ²⁵	Duration ²⁶	Frequency ²⁷	Maximum inter-event period ²⁸	Other requirements
	BF2	Native Fish: NF1, 2, 5, 6, 8 – Recruitment (riverine specialists, generalists) Ecosystem Functions: EF1, 2, 3	September to March	In line with natural ²⁹	5–10 years in 10 (1 every 1–2 years)	2 years	Minimum depth of 0.3 m to allow fish passage
Small fresh	SF1	Native Fish: NF1, 2, 3, 4, 5, 6, 7, 8, 9 – Dispersal / condition (all species) Native Vegetation: NV1 – in-channel Ecosystem Functions: EF1, 2, 3, 4, 5	October to April (but can occur any time)	10 days minimum	Annual (1 per year)	1 year	>20°C for October to April Australian smelt >11°C Minimum depth of 0.5 metres to allow movement of large fish Can follow 2–3 weeks after 'Large Fresh 2' for increased likelihood of successful recruitment of fish, productivity & dispersal Flow ideally up to 0.3 to 0.4 m/s (depending on channel form)
	SF2	Native Fish: NF1, 2, 5, 6, 8 – Spawning (river specialists, generalists) ³⁰ Native Vegetation: NV1 – in-channel Ecosystem Functions: EF2, 3, 4, 5	September to April	14 days minimum	5–10 years in 10 (1 every 1–2 years)	2 years	>20°C; for river blackfish >16°C; for Murray cod September to December >18°C Minimum depth of 0.5 metres to allow movement of large fish Flow ideally up to 0.3 to 0.4 m/s (depending on channel form)

³⁰ Requires the flow to not exceed the large fresh threshold to avoid flushing of nests etc.

Flow component		Environmental objectives	Timing ²⁵	Duration ²⁶	Frequency ²⁷	Maximum inter-event period ²⁸	Other requirements
	SF3	Native Fish: NF1, 2, 3, 4, 5, 6, 7, 8, 9 – Dispersal / condition (all species) Native Vegetation: NV1 - in-channel Ecosystem Functions: EF1, 2, 3, 4, 5, 7 – connectivity between catchments	October to April (but can occur any time)	28 days minimum	Initial flow: Approx. 5–18 months following fish breeding event in Barwon Darling/ large fresh at Wilcannia ³¹ . Follow up flow within 12 months of initial flow to allow further fish movement between catchments	4 years	Minimum depth of 0.5 metres to allow movement of large fish
Large fresh	LF1	Native Fish: NF1, 2, 3, 4, 5, 6, 7, 8, 9 – Dispersal / condition (all species) Native Vegetation: NV1, 3 (in-channel) Function: EF2, 3, 4, 5, 6, 7	Ideally July to September but can occur anytime	5 days minimum	5–10 years in 10 (1 every 1–2 years)	2 years	Flow for pre-spawning condition Minimum depth of 2 metres to cover in-stream features & trigger response from fish Flow ideally 0.3 to 0.4 m/s (depending on channel form)

³¹ Trigger is a demonstrated breeding event in the Barwon Darling with young of year, or, if there has been insufficient monitoring to confirm/deny this, the trigger is a large fresh at Wilcannia. Monitoring results noted in Stuart and Sharpe (2017) *Towards a Southern Connected Basin flow plan: Connecting rivers to recover native fish communities*, indicate that a flow of approximately 10,000 ML/day at Wilcannia coincides with breeding events for golden perch. The 5-month delay following the breeding event is to allow recruits to reach sufficient maturity to move up into the Macquarie.

Flow component		Environmental objectives	Timing ²⁵	Duration ²⁶	Frequency ²⁷	Maximum inter-event period ²⁸	Other requirements
	LF2	Native Fish: NF1, 4, 6, 9 – spawning (flow pulse specialists) Native Vegetation: NV1, 3 (in-channel) Function: EF2, 3, 4, 5, 6, 7	October to April	5 days minimum	3–5 years in 10 years (1 every 2–3 years)	4 years	Rapid rise (comparative to natural rates) >17°C Can be followed by 'Small Fresh 1' for increased likelihood of successful recruitment of fish, productivity & dispersal Minimum depth of 2 metres to cover in-stream features & trigger response from fish Flow ideally 0.3 to 0.4 m/s (depending on channel form)
Overbank / Wetland Small ³²	OB/WS1 (core wetland)	Native Fish: NF1 Native Vegetation: NV2 Waterbirds: WB1, 2, 5 Ecosystem Functions: EF1 – protection of core wetland areas Other Species (Frogs): OS1	Any time	In line with natural (no less than natural) ²⁹	9–10 years in 10	18 months	
	OB/WS2	Native Fish: NF1, 3, 7 – spawning (floodplain specialists) Native Vegetation: NV2, 3, 4a – non-woody wetland	October to April	10 days minimum	5 years in 10 (1 every 2 years)	4 years	>22°C Ideally, recruitment flow 2-4 weeks after spawning flow

³² Flows greyed to denote that discretionary water delivery seeks to avoid floodplain infrastructure and cultivated areas. Otherwise provided by tributary events or flood mitigation zone releases.

Flow component		Environmental objectives	Timing ²⁵	Duration ²⁶	Frequency ²⁷	Maximum inter-event period ²⁸	Other requirements
		vegetation, riparian river red gum Waterbirds: WB1, 2, 5 – habitat Ecosystem Functions: EF2, 3, 4, 5, 6, 7 - connectivity, productivity Other Species (Frogs): OS1 – habitat					
	OB/WS3	Native Fish: NF1, 2, 3, 4, 5, 6, 7, 8, 9 – dispersal & condition (all species) Native Vegetation: NV2, 3 – riparian river red gum communities Ecosystem Functions: EF2, 3, 4, 5, 6, 7 – channel forming, lateral connectivity, productivity Other Species (Frogs): OS1 – habitat	Ideally Sep to Feb but can occur anytime	5 days minimum for fish dispersal In line with natural ²⁹ for riparian river red gum communities	2–3 years in 10 (1 every 3–5 years)	5 years	
	OB/WS4 (Non-woody vegetation zone)	Native Fish: NF1, 2, 3, 4, 5, 6, 7, 8, 9 – dispersal & condition (all species) Native Vegetation: NV2, 3, 4a – Non-woody vegetation in wetlands , river red gum forest Waterbirds: WB1, 2, 3, 5 – habitat Ecosystem Function: EF1, 2, 3, 4, 5, 6, 7 – core wetland	Aug to Mar, with benefits also outside that period including by providing off-season bird	3–8 months for wetlands. Refers to the persistence of standing water, flow can be shorter. For streamside areas, only duration to fill the soil profile,	8–9 years in 10	2 years	Ideally provide slow draw down for shallow muddy edges for bird foraging habitat.


Flow component		Environmental objectives	Timing ²⁵	Duration ²⁶	Frequency ²⁷	Maximum inter-event period ²⁸	Other requirements
		habitats, connectivity to floodplain, productivity Other Species (Frogs): OS1, 2 – small-scale recruitment	foraging habitat	depressions/ billabongs required.			
Overbank / Wetland Medium ³² (river red gum woodland zone)	OB/WM	Native Fish: NF1, 3, 7 – floodplain specialist Native Vegetation: NV2, 4a, 4b, 4e river red gum, lignum Waterbirds: WB1, 2, 3, 4, 5 – habitat & potential small-scale breeding Ecosystem Functions: EF2, 3, 4, 5, 6, 7 – connectivity to floodplain, productivity Other Species (Frogs): OS1, 2 – recruitment	Aug to Mar, with benefits also outside that period including by providing off-season bird foraging habitat	3–6 months for wetlands. Refers to the persistence of standing water, flow can be shorter. For streamside areas, only duration to fill the soil profile, depressions/ billabongs required	5 years in 10	4 years	Ideally provide slow draw down for shallow muddy edges for bird foraging habitat. Ideally have 2 consecutive years of inundation where consolidation of river red gum recruits is desired
Overbank / Wetland Large ³³ (coolibah / black box zone)	OB/WL	Native Fish: NF1, 3, 7 – floodplain specialist Native Vegetation: NV2, 4a, 4b, 4c, 4d, 4e – black box, coolibah & lignum Waterbirds: WB1, 2, 3, 4, 5 – large-scale breeding (colonial & non-colonial) & habitat Ecosystem Functions: EF2, 3, 4, 5, 6, 7 – connectivity to floodplain, productivity	Aug to Mar with benefits also outside that period including by providing off-season bird	3–6 months for wetlands. Refers to the persistence of standing water, flow can be shorter. For streamside areas, only duration sufficient to fill the soil profile,	2 to 3 years in 10	5 years (up to 10 years for outer coolibah/black box areas)	Ideally provide slow draw down for shallow muddy edges for bird foraging habitat.

³³ Greyed to show overbank and wetland flows of this size rely on large tributary events or flood mitigation zone releases.

Flow component		Environmental objectives	Timing ²⁵	Duration ²⁶	Frequency ²⁷	Maximum inter-event period ²⁸	Other requirements
		Other Species (Frogs): OS1, 2 – recruitment	foraging habitat	depressions/ billabongs required.			


4.4 Important flow regime characteristics needed to deliver LTWP objectives

Table 9 Important flow regime characteristics needed to deliver LTWP objectives

Ecological objective	Important flow regime characteristics
 NATIVE FISH OBJECTIVES³⁴	
NF1: No loss of native fish species	<p><u>Cease-to-flow</u>: durations that are not longer than the persistence of water of sufficient volume & quality in key larger river pool refuges for survival of native fish.</p> <p><u>Very low flows (VLF) & baseflows (BF1)</u>: for the survival & maintenance of native fish condition as these flows maintain adequate water quantity & quality (dissolved oxygen, salinity & temperature) in refuge pools.</p> <p><u>Baseflows & small freshes</u>: deep enough along the whole channel to allow fish movement (at least 0.3 m above cease-to-flow for small & moderate bodied fish [Gippel 2013; O'Conner et al. 2015] & 0.5 m for large bodied fish [Fairfull & Witheridge 2003; Gippel 2013; O'Conner et al. 2015]).</p> <p>A <u>baseflow (BF2)</u>: preferably between September & March with an annual or biannual frequency to enhance recruitment outcomes.</p> <p>A <u>small fresh (SF3)</u>: ideally from October to April, but can occur anytime to provide connectivity with the Barwon. This flow should be timed 5 to 18 months after a breeding event in the Barwon-Darling, which is indicated by a large fresh in that system. A follow-up flow 12 months later will allow further recruits from the Barwon to replenish fish populations in the Macquarie-Castlereagh. Duration of 28 days to allow fish to respond to the flow, enter the lower end of the Macquarie-Castlereagh system & work their way up to refuge waterholes higher in the system.</p> <p>A <u>large fresh (LF1)</u>: of at least 5 days duration & occurring ideally between July & September (but can occur at any time) to promote dispersal & pre-spawning condition for all native fish species 5 to 10 years in 10. The large fresh should trigger some primary productivity providing food resources & improving fish condition prior to the spring/summer spawning season. Flow velocities of >0.3 m/s are ideal to trigger fish movement.</p> <p><u>Wetland/overbank flows (OB/WS1)</u>: to maintain core wetlands, including off-channel waterholes. Alternative watering actions (e.g. pumping) may be required to support floodplain habitats under very dry, dry & moderate scenarios to prevent habitat for floodplain specialists from drying out.</p> <p><u>Small overbank & wetland inundating flows (OB/WS3)</u>: ideally from September to February, for at least 5 days & occurring 2 to 3 years in 10 (with a maximum inter-event period of 5 years) is also required to support condition & movement/dispersal outcomes of all native fish groups.</p> <p>Larger flows that inundate off-stream habitat can also promote growth & recruitment through increased floodplain productivity & habitat</p>


³⁴ Important flow regime characteristics for all native fish objectives are based on NSW DPI 2015.

Ecological objective	Important flow regime characteristics
	availability. Larger flows that connect low-lying wetlands provide important habitat to support strong survivorship & growth of juveniles.
NF2: Increase the distribution & abundance of short to moderate-lived generalist native fish species	<p>In addition to the flows listed above for all native fish species, other important aspects of the flow regime for generalists include:</p> <ul style="list-style-type: none"> Regular (ideally annual) spawning & recruitment events for the persistence of short lived species. Spawning can occur independent of flow events, however, <u>small freshes (SF2)</u> enhance spawning. Events should occur during the warmer months of September to April, 5 to 10 years in 10, with a minimum event duration of 14 days for egg development & hatching. Multiple freshes during the spawning season provides flexibility in species response & opportunities for multiple spawning events.
NF3: Increase the distribution & abundance of short to moderate-lived floodplain specialist native fish species	<p>In addition to the flows listed above for all native fish species, other important aspects of the flow regime for floodplain specialists include:</p> <ul style="list-style-type: none"> <u>Overbank & wetland inundating flows (OBWS2)</u> during the warmer months of October to April provide spawning habitat & floodplain productivity benefits to support fish growth. Overbank & wetland flows should inundate floodplain habitats for at least 10 days to allow for egg development & occur at least 5 years in 10, with a maximum inter-event period of 4 years. This period will depend on the persistence of floodplain habitats & time between reconnection to mainstem waterways. Flows should be of a long enough duration to support isolated populations. Water temperatures should be above 22°C. Recruitment is enhanced by subsequent flows events (large fresh, bankfull or overbank & wetland inundating flows) 2–4 weeks after spawning flows. Most floodplain specialist species require spawning & recruitment every 1 to 2 years for population survival.
NF4: Improve population structure for moderate to long-lived flow pulse specialist native fish species	<p>In addition to the flows listed above for all native fish species, other important aspects of the flow regime for flow pulse specialists include:</p> <ul style="list-style-type: none"> A <u>large fresh (LF2)</u> between October to April for a minimum of 5 days for spawning. This is needed 3 to 5 years in 10 with a maximum inter-event period of 4 years. Spawning is triggered by a rapid rise or fall in flow (relative to natural rates) when temperatures are greater than 17°C. In lowland systems, spawning responses are enhanced by substantial flow depths of at least 2 m to cover in-stream features & high flow velocities of greater than 0.3 m/s. Integrity of flow events need to be maintained over long distances (10s to 100s of km) to maximise the capacity for in-stream spawning, downstream dispersal by drifting eggs & larvae & movements by adults & juveniles.
NF5: Improve population structure for moderate to long-lived riverine specialist native fish species	<p>In addition to the flows listed above for all native fish species, other important aspects of the flow regime for riverine specialists include:</p> <ul style="list-style-type: none"> Spawning usually occurs annually, independent of flow events. However, <u>small freshes (SF2)</u> may enhance spawning by promoting ecosystem productivity & inundating additional spawning habitat. Events should occur during the warmer months of September to April, 5 to 10 years in 10. Water temperatures should be >20°C. River blackfish may spawn in lower water temperatures of >16°C & Murray cod in >18°C. Murray cod have a narrower spawning window of September to December. For nesting species (e.g. Murray cod & eel-tailed catfish) preventing rapid drops in water levels (that exceed natural rates of fall) during, & for a minimum of

Ecological objective	Important flow regime characteristics
	<p>14 days after, spawning is important to prevent fish nests from drying.</p> <ul style="list-style-type: none"> Recruitment is enhanced by a secondary flow pulse (large fresh, bankfull or overbank) for dispersal & access to nursery habitat in low-lying wetland habitats. Overall, riverine specialists prefer hydraulically complex flowing streams containing submerged structure (snags & benches) that provides cover & spawning habitat. Flow variability through the delivery of small & large freshes, bankfull & overbank flows enhance the availability of diverse habitat, enhances growth & condition of larvae & juveniles & provides connectivity for dispersal between habitats.
NF6: A 25% increase in abundance of mature (harvestable sized) golden perch & Murray cod	<p>Flow requirements of golden perch (flow pulse specialist) & Murray cod (riverine specialist) are outlined above under NF4 & NF5, respectively.</p> <p><u>Small freshes (SF3)</u> which connect with the Barwon are particularly important for golden perch to provide movement pathways for recruits from breeding events.</p>
NF7: Increase the prevalence &/or expand the population of key short to moderate-lived floodplain specialist native fish species into new areas (within historical range)	<p>Flow requirements of floodplain specialists are outlined for NF3.</p> <p>Expanding populations into new areas will be particularly dependent on dispersal flows, which include large freshes (LF1), & overbank & wetland inundating flows (OB/WS3).</p> <p>Complementary actions such as conservation stocking &/or translocation may be required to support these watering actions. Infrastructure based watering actions (e.g. pumping) may also be required to support floodplain habitats under very dry, dry & moderate scenarios to ensure no loss of species for floodplain specialists (e.g. to prevent wetlands with threatened fish species from drying out).</p>
NF8: Increase the prevalence &/or expand the population of key moderate to long-lived riverine specialist native fish species into new areas (within historical range)	<p>Flow requirements of riverine specialists are outlined for NF5.</p> <p>Expanding populations into new areas will be particularly dependent on dispersal flows, which include large freshes (LF1), overbank & wetland inundating flows (OB/WS3).</p> <p>Complementary actions such as conservation stocking &/or translocation may be required to support these watering actions.</p>
NF9: Increase the prevalence and/or expand the population of key moderate to long-lived flow pulse specialists native fish species into new areas (within historical range)	<p>Flow requirements of flow pulse specialist are outlined for NF4.</p> <p>Expanding populations into new areas will be particularly dependent on dispersal flows, which include large freshes (LF1), overbank & wetland inundating flows (OB/WS3) and flows that connect with the Barwon (SF3).</p>
 NATIVE VEGETATION OBJECTIVES³⁵	
NV1: Maintain the extent & viability of non-woody vegetation communities	<p>Non-woody, inundation tolerant plants occurring on the channel bed, banks, bars & benches require regular wetting & drying to complete life cycles. Variable size & duration of flows including baseflows, variable</p>

³⁵ Important flow regime characteristics for all native vegetation objectives are based on Bowen, S. pers comms, Cassanova 2015, Roberts & Marston 2011, Roberts & Marston 2000, and Rogers & Ralph 2011.

Ecological objective		Important flow regime characteristics
occurring within & closely fringing channels		<p>size freshes & bankfull flows throughout the year will promote diverse communities. Regular inundation will also encourage a dominance of native species over exotic species, as the latter tend to be less tolerant of inundation (Catford et al. 2011). Increased cover of non-woody, inundation tolerant vegetation on banks is likely to stabilise bank material & therefore reduce the risk of excessive bank erosion.</p> <p>Inundation of banks during late winter & early spring by freshes & bankfull flows replenishes soil moisture to promote growth during spring. Prolonged submergence of some amphibious species (e.g. especially if there are continuous high flows) may have detrimental impacts on survival.</p> <p>Small freshes in summer & autumn are important for replenishing soil moisture in river banks to ensure survival & maintenance.</p>
NV2: Maintain the extent & viability of non-woody vegetation communities occurring in wetlands & on floodplains		<p><u>Overbank & wetland inundating flows (OB/WS1, 4):</u> to inundate wetlands for 3 or more months, ideally between August & March, but can occur anytime.</p> <p>The required duration & frequency varies widely by species. Highly water-dependent, amphibious species such as water couch, spike-rush, & cumbungi prefer inundation for 5 to 8 months, 8 to 10 years in 10. The maximum period between events is 2 years. Some amphibious damp species such as floodplain herbs, grasses & sedges that require less frequent (3 to 10 years in 10) & shorter duration (2–4 months) inundation.</p> <p>Other overbank & wetland flows also provide benefit.</p>
NV3: Maintain the extent & maintain or improve the condition of river red gum communities closely fringing river channels		<p><u>Large freshes (LF1, 2):</u> to recharge alluvial aquifers & soil moisture in the riparian zone & maintaining deep rooted vegetation between inundation events.</p> <p>Overbank flows & larger wetland inundating flows (particularly OB/WS3): to inundate the fringing riparian zone.</p> <p>The general condition of riparian vegetation will benefit from inundation or groundwater recharge anytime of the year, with inundation around every 3 years.</p>
NV4: Maintain the extent & maintain or improve the condition of native woodland & shrubland communities on floodplains	River red gum forest & woodland	<p><u>Overbank & wetland inundating flows (OB/WS4 & OB/WM):</u> to inundate vegetation, ideally between 2 & 7 months during September to February. Inundation needs to occur at least 3 years in 10 years, with greater frequency for forested areas.</p> <p>Watering in the year following a maintenance flow will support the survival of seedlings from the previous year in areas where recruitment is desired. Ideally between August to November</p>
	Lignum shrubland	<p><u>Overbank & wetland inundating flows (OB/WM & OB/WL):</u> to maintain condition. For at least 3 months, at a frequency of 3 or more years in 10 & a maximum period between events of 4 years.</p> <p>Regeneration, generally via vegetative expansion requires more frequent inundation (3 to 5 years in 10), for 1 or more months, ideally between September to February.</p>
	Black box woodland	<p><u>Large overbank & wetland inundating flows (OB/WL):</u> to maintain condition, with inundation for 3 to 6 months, at a frequency of 2 to 3 years in 10 & a maximum period between events of 5 years. Regeneration requires watering for 1 to 2 months ideally 5 years in 10.</p>
	Coolibah woodland	<p><u>Large overbank & wetland inundating flows (OB/WL):</u> to maintain condition, with inundation for 2 weeks to 2 months, 1 to 2 years in 10 & a maximum period between events of 10 years. For regeneration a</p>

Ecological objective	Important flow regime characteristics
	duration of 1 to 2 months 5 years in 10 between August & February is required.
 WATERBIRD OBJECTIVES³⁶	
WB1: Maintain the number & type of waterbird species	<p><u>Overbank & wetland inundating flows (OB/WS1, 2, 4; OB/WM; OB/WL):</u> to provide refuge, support feeding & breeding habitat (see WB2, 3, 4) & maintain habitat condition (WB5). Overbank & wetland inundating flows, preferably delivered in spring-summer, that inundate a mosaic of floodplain habitats including non-woody floodplain vegetation, open shallow waterbodies & deep lagoons will provide feeding habitat for a range of waterbird species including open water foragers, herbivores, emergent vegetation dependent species, large waders, wetland generalists & small waders (including migratory shorebird species). Where there is gradual draw-down of habitats over late summer-autumn this can extend feeding habitat available for migratory & resident shorebird species (small waders).</p>
WB2: Increase total waterbird abundance across all functional groups	<p><u>Overbank & wetland inundating flows (OB/WS1, 2, 4; OB/WM; OB/WL):</u> as in WB1 provide seasonal (spring-summer) inundation with gradual draw-down over summer into autumn to provide feeding & breeding habitat (WB3, 4) & maintain the habitat condition (WB5).</p> <p>Where possible to coordinate, these flows should be delivered to compliment events in neighbouring catchments to provide benefits to waterbird populations by providing habitat across a larger area of northern Murray-Darling Basin. Follow-up overbank & wetland inundating flows in years following large breeding events in the Macquarie Marshes & neighbouring catchments in the northern Murray-Darling Basin will also promote the survival of juvenile birds & contribute to increased waterbird populations.</p>
WB3: Increase opportunities for non-colonial waterbird breeding	<p><u>Overbank & wetland inundating flows (OB/WS1, 2, 4; OB/WM; OB/WL):</u> to inundate floodplain habitats for more than 3 months. Spring & summer (preferably September to March) is the ideal season, with opportunistic breeding in autumn & winter.</p> <p>Habitat availability for non-colonial species will increase with increasing magnitude (both extent & duration of inundation) of overbank & wetland inundating flows. Also relies on maintaining (& in some cases improving) the condition of key native vegetation types that provide breeding & foraging habitats (see WB5).</p>
WB4: Increase opportunities for colonial waterbird breeding	<p><u>Overbank & wetland inundating flows (OB/WM; OB/WL):</u> to inundate active colony sites & surrounding foraging habitat for 3 or more months to ensure successful completion of colonial waterbird breeding (from egg laying through to fledging including post-fledgling care) & access to key foraging habitats to enhance breeding success & the survival of young. Ideally September to March.</p> <p>Larger wetland inundation flows (OB/WL) will support larger colonies & a greater number of breeding species with greater benefit to breeding success & increasing total abundance of waterbirds (WB2, WB3). These large overbank events are required on average 2 to 3 years in 10 years, with a maximum inter-event period of 5 years (ideally 4). Medium</p>

³⁶ Important flow regime characteristics for all waterbird objectives are based on Brandis 2010, Brandis & Bino 2016, Rogers & Ralph 2011, and Spencer 2017.

Ecological objective	Important flow regime characteristics
	wetland inundation flows (OB/WM) may support smaller colonies (500 nests) & should ideally occur 5 years in 10.
WB5: Maintain the extent & improve condition of waterbird habitats	<p>Colonial waterbird species are dependent on relatively few sites across the major wetlands of the Murray Darling Basin including known sites in the Macquarie Marshes. These include sites provide nesting habitat consisting of river red gum, river cooba, coolibah, lignum, reedbeds &/or cumbungi.</p> <p><u>Overbank & wetland inundating flows (OB/WS1, 2, 4: OB/WM: OB/WL)</u> are needed to maintain the extent & condition of these vegetation communities in these discrete wetland sites. This ensures that sites are in event-ready condition when medium & large events (<u>OB/WM: OB/WL</u>) initiate colonial waterbird breeding events. These flows, particularly the smaller overbank & wetland inundating flows (OB/WS1,2,4), will also support a broader range of foraging habitats in the Macquarie Marshes, including spike-rush sedgeland, marsh grasslands, lignum shrublands & open lagoons. The required duration & frequency of overbank flows to support these vegetation types are outlined under the native vegetation objectives.</p>




PRIORITY ECOSYSTEM FUNCTIONS OBJECTIVES³⁷

EF1: Provide & protect a diversity of refugia across the landscape	<p><u>Cease-to-flow</u>: durations that are not longer than the persistence of water of sufficient volume & quality in key larger river pool refuges.</p> <p><u>Very low flows (VLF) & baseflows (BF1)</u>: to maintain adequate water quantity & quality (dissolved oxygen, salinity & temperature) in refuge pools. Need to be of sufficient magnitude to prevent stratification of pools that can lead to de-oxygenation of the water column. They are required every year at a frequency that is no less than natural, & are especially important during dry times.</p> <p><u>Small freshes</u>: when restarting flows after a cease-to-flow event, larger magnitude flows such as these may be required to prevent detrimental water quality outcomes (as poor-quality water from the bottom of pools is mixed through the water column).</p> <p><u>Wetland/overbank flows (OB/WS1)</u>: To maintain core wetlands, including off-channel waterholes. Alternative watering actions (e.g. pumping) may be required to support floodplain habitats under very dry, dry & moderate scenarios to prevent habitat for floodplain specialists from drying out.</p>
EF2: Create quality instream & floodplain habitat	<p>The full range of in-channel & overbank flow types are required to maintain quality instream & floodplain habitat. Variable in-channel flows (baseflows – bankfull flows) will provide a diversity of physical & hydraulic habitats. With increasing magnitude of flows, greater areas of the channel are inundated (e.g. benches, bars, snags & banks at different elevations in the channel). Baseflows & small freshes provide areas of slackwater (slow flowing) habitat, while large freshes provide deeper & faster flowing habitats. Small & large freshes are important for flushing fine sediment from pools, de-stratifying pools & maintaining geomorphic features such as benches & bars. Bankfull flows are important for geomorphic maintenance of all channel features.</p> <p>To protect banks from excessive erosion it is important to maintain rates of fall that do not exceed natural rates of fall for all regulated deliveries.</p>

³⁷ Important flow regime characteristics for all priority ecosystem function objectives are based on Alluvium 2010.

Ecological objective	Important flow regime characteristics
	<p>Slow rates of fall allow water to drain from the bank slowly, preventing mass failure of the banks. Maintaining slow rates of fall is particularly important when flows are in the lower third of the channel, to protect the 'toe' of the bank, which supports the rest of the bank above.</p> <p>Bank notching can be avoided by varying flows (avoiding holding flows constant for too many consecutive days) & targeting different peak heights for freshes.</p> <p><u>Wetland/overbank flows (OB/WS1-4; OB/WM; OB/WL):</u> to provide floodplain & wetland habitat for native fish, waterbirds & other aquatic fauna.</p>
EF3a: Provide movement & dispersal opportunities within catchments for water-dependent biota to complete lifecycles	<p>Providing longitudinal connectivity is critical for migration, recolonisation following disturbance events, allowing species to cross shallow areas, & dispersal of larvae to downstream habitats. In-channel flows of adequate depth & duration (baseflows & freshes) are important to allow for the movement of aquatic & riparian fauna & flora along rivers & creeks. For example, flows of at least 0.3 m are needed to allow medium sized native fish to move along a channel. Physical barriers, such as dams & weirs, have introduced additional barriers throughout the Macquarie-Castlereagh, making large freshes & occasionally small overbank flows important for overcoming these man-made structures where fishways are not present.</p> <p><u>Wetland/overbank flows (OB/WS1-4; OB/WM; OB/WL):</u> to provide lateral connectivity to floodplain & wetland habitat.</p>
EF3b: Provide movement & dispersal opportunities between catchments for water-dependent biota to complete lifecycles	<p><u>Freshes & baseflows:</u> to connect the Lower Macquarie/Marthaguy, Marra, Castlereagh & Bogan to the Barwon are important to allow the movement of fish & other biota between the catchments & allow re-colonisation & the replenishment of populations.</p> <p>Particularly <u>small fresh (SF3)</u>: see description of this flow under NF1.</p>
EF4: Support instream & floodplain productivity	<p><u>Freshes:</u> to flush organic matter from stream banks & anabranches & drive small pulses of productivity.</p> <p><u>Wetland/overbank flows (OB/WS1-4; OB/WM; OB/WL):</u> to inundate the floodplain & supporting large scale productivity, which in turns drives aquatic food webs both on the floodplain & in-stream. Primary productivity includes growth of algae, macrophyte, biofilms & phytoplankton, which in turn drives secondary productivity (zooplankton, macroinvertebrates, fish larvae etc.).</p>
EF5: Support mobilisation & transport of sediment, carbon & nutrients along channels, between channels & floodplains, & between catchments	<p><u>Freshes:</u> for mobilising organic matter & sediment from in-channel surfaces (e.g. leaf litter that has accumulated on bars, benches & banks during low flows). This material is transported downstream or deposited in other parts of the channel where it is utilised, in the case of nutrients & carbon, to drive primary productivity, or in the case of sediment, for channel maintenance (e.g. to replenish banks & benches).</p> <p><u>Wetland/overbank flows (OB/WS1-4; OB/WM; OB/WL):</u> for transferring nutrients & carbon from the floodplain to the channel.</p>
EF6: Support groundwater conditions to sustain groundwater-dependent biota	<p><u>Large freshes (LF1, 2) & wetland/overbank flows (OB/WS1-4; OB/WM; OB/WL):</u> to contribute to recharging shallow groundwater aquifers in areas where there is a surface-groundwater connection. This recharge can reduce the salinity of shallow aquifers & raise water tables, providing critical soil moisture for deep-rooted vegetation in the riparian zone & on low-lying floodplains.</p>

Ecological objective	Important flow regime characteristics
EF7: Increase the contribution of flows into the Murray & Barwon-Darling from tributaries	<p>Flows from the Macquarie-Castlereagh to provide movement & dispersal opportunities between catchments (see EF3b) will also contribute to important EWRs in the Barwon-Darling WRPA.</p> <p>Protecting <u>overbank flows (OB/WS2-4; OB/WM; OB/WL)</u> will provide important flows & transfer nutrients & carbon from floodplains in the Macquarie-Castlereagh to the Barwon River.</p>
 OTHER SPECIES ³⁸	
OS1: Maintain species richness & distribution of flow-dependent frog communities	<p>In addition to actions which allow breeding (OS2) the flows below are important for survival & to maintain frog condition.</p> <p><u>Cease-to-flow</u>: durations that are not longer than the persistence of water of sufficient volume & quality in key larger river pool refuges.</p> <p><u>Very low flows (VLF) & baseflows (BF1)</u>: to maintain adequate water quantity & quality (dissolved oxygen, salinity & temperature) in refuge pools.</p> <p><u>Wetland/overbank flows</u>: Flows to maintain core wetlands, including off-channel waterholes (OB/WS1) for refuge. Larger flows (OB/WS2-4, OBWM, OBWL) to maintain frog condition & habitat & allow dispersal.</p>
OS2: Maintain successful breeding opportunities for flow-dependent frog species	<p><u>Wetland/overbank flows (OB/WS4, OB/WM, OB/WL)</u>: to provide opportunities for breeding & recruitment (i.e. laying eggs & tadpole metamorphosis). Ideally every 1 to 2 years for 6 or more months (with a minimum of 4 months). Spring-summer breeders require flows ideally from October to March, while species with more flexible breeding are likely to benefit from flows arriving between July & April. A gradual rise & fall is likely to improve recruitment outcomes.</p>

³⁸ Important flow regime characteristics from J. Spencer and J. Ocock (OEH, pers. comms. 2018)

4.5 Changes to the flow regime

The flow regime in the Macquarie-Castlereagh WRPA has changed due to regulation and development in the catchment. The degree of change varies depending on the location within the catchment. This section:

- briefly describes key differences between the pre-development and post development flow regimes
- notes aspects of the current flow regime which impact on achieving EWRs in some PUs. This assists in identifying risks and framing priority management actions.



Figure 9 Burrendong Dam offtake tower with cold water curtain. Photo: Tim Hosking/OEH

Areas that are regulated or affected by regulated water

Differences in the flow regime between pre-development and post-development conditions are discussed below based on observations (where available) and modelled scenarios.

Observations assess changes in recorded flows prior to the 1967 completion of Burrendong Dam ('pre-development') compared to flows recorded after Windamere Dam's 1983 completion ('post-development'). However, note this may understate the effect of development because there was development prior to 1967 and further development occurred after 1984. Note also that different time periods are compared, so there will be differences due to natural variations between years, not just due to water use development.

Modelled comparisons use modelled flows under a 'without development' scenario (without dams or extractions) and under a 'current conditions' scenario and compare these. NSW DOI-W undertook the modelling and analysis and presented it in the *Risk Assessment for the Macquarie-Castlereagh Water Resource Plan Area* (DOI-W in prep). Note the DOI-W comparisons were done prior to the development of the EWRs outlined in this plan (Section 4.3). They analyse baseflows, freshes and high/overbank flows³⁹ based on the frequency of occurrence, and these flows may not line up precisely with the baseflows, freshes and overbank flow levels in the EWRs of this LTWP.

Cudgegong River between Windamere Dam and Burrendong

Modelling by DOI-Water (in prep) shows that with development there has been a greater than 50% increase in baseflows at the expense of high and overbank flows. High and overbank flows have decreased in size by more than 50% at Rocky Waterhole, but this effect is lessened downstream at Yamble Bridge where the reduction is less than 50%, due to tributary contributions between Rocky Waterhole and Yamble Bridge.

Modelling shows a mixed effect on freshes, with an increase at Rocky Waterhole but a decrease at Yamble Bridge. Further work will be needed to determine the causes of this.

The seasonality of flows has also changed with median flows now higher in summer and lower in winter.

Macquarie River from Burrendong Dam to Warren Weir

As with the regulated Cudgegong, modelling by DOI-W (in prep) shows that with development there has been a greater than 50% increase in baseflows at the expense of high and overbank flows. High and overbank flows have generally decreased in size by more than 50% down to Gin Gin, with less substantial decreases at Warren. The smaller size of the decrease downstream at Warren may be due to the natural reduction in 'peakiness' in flows lower in the catchment. This means that larger flows do not always show up in significantly higher peak flows, but in marginally higher peaks that last longer.

Modelling indicates that freshes have increased. Analysis of observed flows shows some differences between small freshes, which have generally increased at Dubbo and Warren Weir since development, and large freshes, which have decreased in frequency. The number of years LTWP large fresh EWRs (LF1 and LF2) are met has decreased by more than 30% at Dubbo. At Warren Weir the decrease is less pronounced, but still 11% for LF1 and 32% for LF2. It is also worth noting that although small freshes appear to have increased in frequency, these freshes now generally come from Burrendong Dam deliveries. These will have a different ecological value to natural rainfall/runoff induced freshes. These have different rates of rise and fall and durations and contain more organic matter and different sediment/chemical signatures. All of these aspects can serve as cues for aquatic biota to respond to.

The seasonality of flows has also changed with median flows now far higher in summer and lower in winter.

Macquarie Marshes flows

By the time flows reach the Macquarie Marshes, it appears that baseflows are no longer higher than pre-development times. Modelling by DOI-W (in prep) indicates a distinct decrease (greater than 50%). However, observations at Oxley show a less distinct reduction with the number of days per year of baseflow dropping by only 10% from 1943–1966 to 1984–2017.

³⁹ Flows with an average return period of 1.5 years or more, and termed 'high and infrequent flows' in DOI-W (in prep).

DOI-W (in prep) modelling shows that the size of high and overbank flows has dropped, but by less than 20%. There are no significant tributaries between Warren and the Marshes, so the limited magnitude of change seen in the Marshes compared to upstream areas is likely due to the reduction in the 'peakiness' of flows lower in the catchment.

Modelling by DOI-W (in prep) finds that freshes have increased by more than 50% on the Macquarie River at Marebone and Oxley. However, the levels used to measure freshes are different to those used in the LTWP, and analysis of observed flows at Oxley show different results for the small and large freshes used in the LTWP. These show a low (less than 20%) reduction in the frequency of years the LTWP small and larger freshes EWRs (SF1, SF2, LF1, LF2) occur from 1943-1966 compared to 1984-2017.

The seasonality of flows has changed with median flows now far lower in June to September. Unlike in the upstream regulated river that had increases in summer, this is not seen to the same extent in the Marshes, most likely because the higher summer flows seen upstream have been for irrigation deliveries and have been extracted above the Marshes.

Macquarie Marshes inundation events

DOI-W (in prep) did not analyse inundation events in the Marshes, however there is other information about these events which is useful for water management.

Thomas et al. (in prep) calculated the area of the Marshes covered by inundation events (flow scenarios) of different sizes in the Marshes using satellite imagery from 1988–2013. The size of the flow scenarios analysed is at Table 10 and the extent of the Marshes they covered is mapped at Figure 11.

Table 10 Predicted inundated areas for specific flow scenarios

Flow Scenario (at the combined Macquarie River and Marebone Break below Marebone Weir gauges)		Predicted Inundated Area (nearest 1,000 ha)
Volume (over number of days)	Inundation Zone colour	Mean area
30,000 ML (90 days)	Blue	8,000
60,000 ML (90 days)	Purple	14,000
100,000 ML (90 days)	Pink	21,000
250,000 ML (120 days)	Red	45,000
400,000 ML (150 days)	Orange	65,000
700,000 ML (180 days)	Green	103,000
1,000,000 ML (280 days)	Yellow	138,000

OEH has analysed available observed data to identify the largest flow scenario that occurred in each year since gauging began in 1986 (presented at Figure 11).

From the sequence of events seen in Figure 11, one of the key observations is the run of dry years from 2006/07 to 2008/09. Successive dry years such as this are likely to negatively affect the condition of the core Marshes and result in minimal refuge habitat availability during those years. These results highlight the importance of reducing the length of such dry runs. This can be done by carrying water over from wet and moderate years to service the needs of dry years. The strategy used for carryover will be crucial and it will need to balance the potential benefits and opportunity costs of carrying over different volumes of water in any given year. Further work on this strategy is identified in Table 19. The effectiveness of carryover is reliant on carryover being available in the resource assessment, so it is important that environmental water continues to maintain the same rights to access of carried over water as other water users.




Figure 10 Areas of the inundation in the Macquarie Marshes based on flow scenarios of different sizes listed in Table 10 from Thomas et al. (in prep)




Figure 11 Occurrence of events in Macquarie Marshes, based on flow scenarios of different sizes listed in Table 10 from Thomas et al. (in prep).

Note the yellow zone is not shown and some larger green events would have been in the yellow zone.

Lower Macquarie River

As the lowest reach of the Macquarie, this stretch is the most heavily impacted by the cumulative effects of river regulation and extraction (including floodplain harvesting, unregulated licenced extraction and regulated extraction). Analysis of modelling by DOI-W (in prep) shows all flow classes (baseflows, freshes, high and overbank flows) have reduced by more than 20%.

Analysis of observed flows at the Macquarie River Bell's Bridge (Carinda) gauge also shows similar reductions in flows between 1938-1966 and 1984-2017. Note there is further extraction downstream of this gauge, so the effect is likely to be even greater by the time flows reach the Barwon River.

Concerns to be highlighted are the:

- reduction in small and large freshes will impact on the connectivity to the Barwon for fish and other organisms.
- reduction in freshes and overbank flows, because these provide important flow volume contributions to the Darling-Barwon system.

Analysis of observed flows shows there has also been a change in the pattern of events between 1938-1966 and 1984-2017, with events now being shorter but more frequent. This may be because larger events are being broken up into several smaller ones through regulation or extraction during events.

Lower Marthaguy Creek

DOI-W's (in prep) analysis of modelled flows shows that base flows and freshes have increased by over 50% at Carinda. This is downstream of the confluence with the Terrigal system, so contributions from deliveries to the Gum Cowal/Terrigal system may be providing increases here. However, analysis of observations at the Marthaguy Creek Carinda gauge show minimal change in the average number of days of baseflow from 1944–1966 to 1984–2017 (decrease from 116 to 113 days per year for flows reaching 20 ML/day). Analysis of observations for small and large freshes also show a reduction in the number of years that flows of the required duration to meet the EWR requirements (see Table 8). The differences between the modelled and observed assessments may be due differences in flow rates being assessed.

Modelling by DOI-W (in prep) also shows that high and overbank flows with average return periods of 1.5 and 2.5 years have decreased in size by more than 20%, with decreases also seen in larger 1-in-5 year floods, although the reduction is less than 20%. The lower Marthaguy Creek provides flows and connectivity to the Barwon River (via the Lower Macquarie) and so reductions in flows from the Marthaguy also have wider implications.

Marra Creek

Key gauges in this creek (at Carinda Road and at Billybingbone Bridge) only begin in the 1980's so there is no pre-development observed data available. This makes it difficult to determine the type and degree of effect in this system. However, because high flows in Macquarie River have reduced it is possible to make inferences from this. The Marra Creek's natural off-take channel (or 'break') starts upstream of Marebone Weir and has a commence-to-flow of about 3200 ML/day (Torrible et al. 2011). Modelling results presented by Torrible et al. (2011) support this, showing the pre-development one-in-two year flood now occurs only one-in-five years. As with the Lower Macquarie River and Marthaguy Creek, the Marra Creek provides flows and connectivity to the Barwon River, meaning the impacts of flow reduction indicated by this will extend beyond the creek itself. Due to deficiencies in the data it is recommended that further work be undertaken to determine the optimal flow regime for this creek, building on analysis undertaken for this LTWP (see Table 19).

Distributary (or effluent) creeks: Gunningbar, Duck and Crooked creeks.

As with Marra Creek, key gauges on these creeks only begin with continuous records in the 1980's so there is no pre-development observed data available. Analysis of modelled data by DOI-W (in prep) shows zero flow periods in Gunningbar, Duck and Upper Crooked creeks have actually become rarer and baseflows have increased by over 50%. However, freshes and overbank flows have decreased by over 50% in all these creeks.

Torrible et al. (2011) reported on the environmental effects of the flow regime in some of these creeks. The effect of reduced high flows was seen in the Talga Wetlands/Overflow of the Lower Crooked Creek, which is reported to have supported significant bird breeding events in wetter times and have been dominated by marsh club rush (where this forms stands or 'sedgelands', it is a critically endangered community under the *Biodiversity Conservation Act 2016*). This is now in a degraded state. Torrible et al. (2011) noted the improvement of the condition of the low-lying floodplains of Lower Crooked Creek as a priority *where feasible*. Because the feasibility of water delivery in this area currently considered low, further work is needed in this area (see Table 19).

In regard to baseflows, Torrible et al. (2011) and Barma et al. (2011) recommended for Gunningbar Creek that flow permanency be reduced to provide a more variable flow pattern and a more natural wetting-drying regime. Barma et al. (2011) recommended the same for Duck Creek. Section 7 of this LTWP notes this as an area for further work (Table 19).

Unregulated areas

Unregulated licenced extractions and extraction for basic landholder rights are the main factors influencing the flow regime in unregulated areas. They can increase cease-to-flow periods, decrease pool persistence under dry conditions and reduce low flows. The frequency of small fresh events can also be decreased. While total licenced extractions in unregulated areas is generally lower than in regulated areas, changes to the return frequency of flow components can impact on the character and persistence of flow-dependent ecological communities. Note that in some 'unregulated' areas, particularly near larger population centres, town water supply dams may also significantly affect flows.

Ecological values are high in many of these unregulated PUs. Flows in these systems are entirely dependent on natural rainfall events and are not manageable except by some small-scale structures such as small weirs and in-stream (licenced) dams. The main tool available to maintaining or improving flow regimes is through the rules in the WSP, which define when water can be taken, the limits to extraction and on trading into management zones. Management recommendations to support important flows in unregulated PUs are described in Part B of the LTWP.

5. Risks, constraints and strategies

This LTWP is focussed on delivering environmental outcomes in a heavily modified landscape. There are many factors that could potentially impact on how the plan is implemented, and how the environment responds to management actions.

The *Risk Assessment for the Macquarie-Castlereagh Water Resource Plan Area* (DOI-W in prep) identifies risks to areas of conservation value, based on hydrological change within sub-catchments, and outlines how the Water Resource Plan may mitigate those risks. This section complements the Macquarie-Castlereagh risk assessment and addresses the specific risks and constraints that may arise in the implementation of the LTWP.

The risks and constraints outlined in this LTWP are those that affect our capacity to:

- meet the EWRs of environmental assets and functions (Table 11)
- achieve the environmental objectives of the LTWP (Table 12).

This risk assessment has assisted with identification of appropriate investment opportunities for improving the likelihood that EWRs can be achieved (Table 19).




Figure 12 Sunset over Macquarie Marshes. Photo John Spencer/OEH

5.1 Risks and constraints to meeting EWRs in the Macquarie-Castlereagh WRP

Table 11 Water management related risks and constraints to meeting objectives and potential strategies for managing them

Risk	Description	Potential management strategies	Responsible organisations
Insufficient water for the environment to meet LTWP objectives	Availability (total volumes, season, flow rate) of water for the environment does not meet environmental demand.	Regulated (or affected by regulated water) areas	
		Continue to enable adaptive water management through review of WRP outcomes against LTWP objectives in 5 years (2024).	OEH, DOI-W
		Ensure future water management policy & WRP changes that affect the availability of water for environmental outcomes considers the impact on progress toward achieving environmental objectives.	DOI-W, MDBA
		Continue to improve monitoring, research & assessment methods to inform decision making.	OEH, CEWO, Fisheries, DOI-W, WaterNSW
		Investigate options for the strategic delivery of irrigation orders to mimic natural flow events (requires interagency discussion).	WaterNSW, OEH & DOI-W
		Unregulated areas	
		Maintain & where necessary implement new rules restricting trade into water sources with high or medium risks (as defined by the WRP Risk Assessment, DOI-W (in prep)).	DOI-W
		Review low-flow access rules where in-channel flows have been highly impacted since development (as defined by the WRP Risk Assessment, DOI-W (in prep)).	DOI-W
		Develop & implement the floodplain harvesting rules of the water sharing plan & WRP for the Macquarie-Castlereagh WRP.	DOI-W, NRAR
		Ensure the plan recognises the impact of such harvesting on downstream areas & on connectivity & contributions to the Barwon system. Ensure compliance with those floodplain harvesting rules.	
		Consider trade out of high-risk areas as a mechanism to ensure that sufficient water is retained for the environment.	DOI-W

Risk	Description	Potential management strategies	Responsible organisations
Extraction & diversion of environmental water for purposes that are not consistent with LTWP objectives	<p>This includes: Currently legal extraction of environmental water orders below the gauged delivery point (e.g. Marebone Weir), reducing benefits to downstream assets & end-of-system flows Unauthorised extraction or diversion of flows. Diversion of water from targeted environmental flows due to unsympathetic works approval rules Increasing substitution of stock & domestic supply with environmental water.</p>	Refer to the Natural Resources Access Regulator & MDBA water compliance policies & strategies	NRAR, MDBA
		Compliance monitoring	NRAR, MDBA
		Implement protection of environmental flows as recommended by the Matthews report (2017a, b).	DOI-W & WaterNSW
		Communicating the whole-of-system management approach to help improve understanding of the importance of protecting environmental flows beyond the order point.	OEH & DOI-W
		In unregulated PUs implement management strategies outlined in Section 6.2 <i>Protection of ecologically important flow components in unregulated river reaches</i>	DOI-W & WaterNSW
		Review problematic structures for their consistency with environmental water objectives. Where they are inconsistent, change WSP rules & licence conditions to properly reflect their purpose & water source.	DOI-W & WaterNSW
		Review S&D supply in some areas – Milmiland Creek, Gum Cowal, Lower Macquarie River & Bulgeraga Creek to ensure managed environmental water is not being substituted for recognised supply from the WSP socialised pool – representing incremental erosion of PEW volumes & reducing the effectiveness of HEW outcomes.	DOI-W & WaterNSW
Floodplain structures & barriers preventing flows that would meet overbank & wetland inundation EWRs	Construction of structures (e.g. levees, diversion channels, sediment blockage of culverts) that causes barriers to flows to wetlands & ecological-important floodplain areas.	<p>Implement the <i>Floodplain Management Plan for the Macquarie Valley Floodplain</i> (OEH 2018, currently in draft)</p> <p>Monitor compliance of infrastructure with the Floodplain Management Plan for the Macquarie Valley Floodplain.</p> <p>Note that any flow impeding infrastructure in & directly upstream of the Marshes area will have particularly detrimental effects on the Eastern Marshes.</p>	DOI-W, NRAR
Insufficient channel or dam valve capacity to deliver flow requirements	Due to Burrendong Dam valve configuration, there is limited capacity during some	Investigate issues & improve delivery of environmental water at times of high channel use.	OEH & WaterNSW
		Environmental & irrigation entitlements should maintain the same delivery rights.	OEH & WaterNSW

Risk	Description	Potential management strategies	Responsible organisations
	seasons to deliver water.	Develop formal supply sharing arrangements to provide a mechanism for managing the delivery of water when demands regularly exceed channel sharing capacity.	OEH & DOI-W
	Delivery to some areas (e.g. Crooked Creek, Marra Creek) is constrained by the capacity of inflow channels.	Support the ongoing integration of environmental water planning strategies into river operations to achieve environmental outcomes with all water.	OEH & WaterNSW
	Cudgegong River flows are restricted to around 1,500 ML/day by the Rocky Waterhole bridge	Support proposed upgrade of Cudgegong River crossings to allow higher flows (see Table 19).	WaterNSW
Rate & timing of irrigation releases not optimised for environmental outcomes	Consumptive water is typically released from Burrendong Dam as efficiently as possible. There are opportunities to improve the environmental performance of river operations.	Optimising water releases for multiple benefits	OEH, WaterNSW & DOI-W
		Gradual declines in water level after flow events will benefit the ecology & bank stability of the river. This could be achieved through changes to river operations or the use of environmental water to assist.	
		Strategic delivery of irrigation orders to mimic natural flow events	
Inappropriate commence & cease-to-pump rules in unregulated catchment impacting on some flow classes & EWRs	Visible flow can be an ambiguous trigger for pump rules & compliance is difficult to enforce. There are not enough gauges in some areas of the unregulated WRPAs to support these rules.	Investigate whether the risk associated with the reduced frequency of freshes during late spring & summer can be mitigated by delivering bulk water in patterns that mimic natural flow conditions. The ability to implement this strategy will vary between years & seasons & must be consistent with the need for efficient & timely water delivery.	OEH, DOI-W, WaterNSW & CAG
		Investigate improved metering of pumps. Note the Matthews (2017a, b) reports recommended universal water metering.	DOI-W
Inappropriate pool drawdown rules impacting on drought refuge EWRs	Pumping from pools during dry periods impacts on valuable drought refuge. Limited gauging & metering makes	Investigate better stream gauging to help licence holders & compliance officers determine stream flow (Table 19).	DOI-W
		Ongoing & improved monitoring & evaluation to inform revision of pool drawdown rules in the <i>Macquarie-Castlereagh Water Sharing Plan</i> to ensure they mitigate impact on high-value refuge sites.	OEH, Fisheries & DOI-W

Risk	Description	Potential management strategies	Responsible organisations
	compliance monitoring difficult.		
Impacts to floodplain infrastructure & operations constraining meeting some EWRs	The risk of inundation of cultivated land, disrupted access & inundation of stock & infrastructure can cause impacts to businesses & residences.	Improve stakeholder education & resources to increase understanding of floodplain inundation patterns.	OEH, CEWO, DOI-W, WaterNSW, LLS, CAG, & Landholders
		Implementation of the Floodplain Management Plan for the Macquarie Valley Floodplain (OEH 2018, currently in draft).	DOI-W
		Consider inundation risks when planning water deliveries.	OEH & CEWO
		Communicate with landholders that may be affected by flows about intended water deliveries.	OEH
		Monitor natural events & environmental deliveries to determine the risk of inundation under a range of flow rates.	OEH & CEWO
		Provide regular updated information for landholders during higher flows.	WaterNSW
		Investigate options for relaxing constraints to delivery of environmental flows, particularly to the Macquarie Marshes. See Table 19.	DOI-W & OEH
Stream channelisation compromising meeting overbank flow EWRs	Stream channel erosion can deepen the channel & prevent flows from spilling out onto the floodplain	Consider channel restoration works to reinstate appropriate channel profile, noting potential impacts from altered hydrology. See Table 19.	OEH

5.2 Non-flow related risks and constraints to meeting LTWP objectives

The risks and constraints to meeting the ecological objectives include non-flow-related external factors that could potentially impact on achieving the objectives outlined in this plan (Table 12). While managing these risks and constraints is outside the scope of this LTWP, they have been included to draw attention to their influence on river and wetland health, and to highlight the importance of linking this LTWP with broader natural resource management.

Table 12 Non-flow management related risks and constraints to meeting LTWP ecological objectives

Risk	Description	Potential management strategies	Responsible organisations
Poor water quality, including hypoxic blackwater and thermal pollution impacts affecting LTWP fish & functions objectives	<p>Water quality affects the ecology & survival of aquatic organisms. Hypoxic blackwater events can occur with the release of water after dry or low-flow periods. This can occur from the build-up of organic material in channels & on floodplains. Can lead to low-dissolved oxygen levels & potential fish kills. Thermal pollution can occur when curtain is off-line or multi-level offtake is not operated optimally for temperature outcomes in instances when algal blooms are present</p>	Implement recommendations detailed in the <i>Water Quality Salinity Management Plan</i> (DOI-W, in prep)	DOI-W, OEH
		Manage salinity in accordance with the <i>Basin Salinity Management 2030 Strategy</i> (MDB Ministerial Council 2015)	DOI-W
		Consider potential for benefits to water quality when managing water to meet environmental objectives.	OEH, CEWO
		Implement land management strategies to improve water quality.	LLS with Landholders, Landcare, OEH, WaterNSW, DOI-W & other community groups
		Consider risks when delivering environmental flows during high-risk periods, such as warm weather in late spring & summer.	OEH & CEWO
		Provide flow regimes that avoid extended dry or very low-flow periods.	Water NSW, OEH & DOI-W
		Restart rivers with flow rates that reduce the risk of hypoxic blackwater, informed by water quality monitoring.	OEH & WaterNSW
		Operation and maintenance protocol for Burrendong curtain developed & implemented Strategy for Windamere multi-level offtake operation implemented	WaterNSW
Native vegetation clearing impacting on LTWP vegetation & waterbird habitat objectives	<p>Native vegetation clearing has direct impacts on LTWP vegetation objectives & the availability of waterbird habitat. Changes to riparian vegetation can impact on water quality, stream erosion & instream habitat.</p>	Work with relevant departments & organisations to identify & protect core wetland vegetation communities using legislation & native vegetation planning processes.	OEH, LLS & DPI
		Map & identify riparian & aquatic habitat condition to inform development of formal agreements in a unified strategy. Prioritise reaches for management in partnership with LLS & landholders. See Table 19.	OEH, Fisheries & LLS
Impacts of unmanaged total grazing pressure & stock access to waterways	If not managed carefully, grazing pressure from domestic & native herbivores & access of	Map & identify riparian & aquatic habitat condition to inform development of formal agreements in a unified strategy. Prioritise reaches for management in partnership with LLS & landholders. See Table 19.	OEH, Fisheries & LLS

Risk	Description	Potential management strategies	Responsible organisations
impacting on LTWP vegetation targets	stock to riverbanks can: <ul style="list-style-type: none"> • reduce native vegetation cover which allows weeds to establish • reduce streambank stability • damage important instream habitat • reduce water quality. 	Implement grazing strategies that protect & restore wetland vegetation as per Holmes <i>et al.</i> (2009) www.dpi.nsw.gov.au/agriculture/pastures-and-rangelands/establishment-mgmt/grazing-management2/guidelines-gwydir-macquarie .	LLS, DPI & Landholders
		Investigate incentives to improve management of wetlands on private land.	LLS, DPI, BCT
		Manage the abundance the abundance of native herbivores to control total grazing pressure.	OEH National Parks & Kangaroo Management, landholders & LLS
Spread of pest plant species impacting on achieving LTWP vegetation objectives	There is potential for environmental water to promote the growth & spread aquatic & terrestrial weed species including lippia, Noogoora burr, tobacco weed.	Map & identify riparian & aquatic habitat condition to inform development of formal agreements in a unified strategy. Prioritise reaches for management in partnership with LLS & landholders. See Table 19.	OEH & LLS
		Maintain existing weed control programs.	OEH National Parks, LLS & local government
		Inundate wetlands for enough time to favour native wetland species growth & reduce the extent of terrestrial weed species like lippia.	OEH & CEWO
		Monitor for pest species, including for potential new pests.	DPI, OEH (including National Parks), LLS, landholders & local government
Spread of pest animal species impacting on achieving LTWP vegetation, fish, frog & waterbird objectives	The current flow regime, including environmental water supports populations of invasive animals. These populations reduce the benefit of environmental water on native species.	Support recommendations in pest species management plans, with implementation of control programs such as those for: <ul style="list-style-type: none"> • carp (see NSW I&I 2010), including the cyprinid herpesvirus-3 if recommended by the National Carp Control Program (FRDC in prep) • other invasive fish such as redfin & gambusia (competition with native fish & predation) • pigs (vegetation, predation impacts) • goats (vegetation impacts) • foxes & cats (predation impacts). 	OEH National Parks, Fisheries, WaterNSW & OEH

Risk	Description	Potential management strategies	Responsible organisations
		Implement proposals in <i>Fish for the Future: Action in the Northern Basin—NSW proposal for Northern Basin Toolkit measures to promote native fish health.</i> (NSW Fisheries, 2017)	Fisheries
		Monitor for pest species, including for potential new pests (e.g. Tilapia).	Fisheries, OEH (including National Parks), LLS, landholders & local government
Problematic erosion & sedimentation may impact upon various LTWP objectives	Erosion (both natural & accelerated) & sedimentation may result in a variety of changes that affect the LTWP targets, including: <ul style="list-style-type: none"> vegetation (inundation extent change, changes in streams) waterbird habitats water quality longitudinal & lateral connections 	Support variable flows & ecologically desirable flow recession rates in river operations to reduce bank slumping.	OEH & WaterNSW
		Map & prioritise riparian habitat & erosion points for rehabilitation at the catchment scale, with a commitment to manage risk & monitor outcomes. See Table 19.	OEH, LLS, Fisheries & WaterNSW
		Manage environmental waters to mimic pre-development flow patterns & variability (where possible).	OEH, DOI-W, WaterNSW, CEWO
		Investigate ongoing erosion control works in high-risk areas, such as the Southern Marshes & key breaks	OEH, WaterNSW, DOI-W
		Treat known erosion risk areas e.g. Bell River confluence, southern Marshes	LLS, DOI-W, Councils, OEH landholders
Current & future instream barriers & structures impacting connectivity related LTWP objectives	There are 238 major barriers including weirs, regulators, & road crossings identified in the Macquarie-Castlereagh. These impede natural flow & connectivity, significantly impacting fish populations.	Remove priority illegal & unauthorised barriers.	DOI-W, Fisheries
		Implement NSW Department of Primary Industries, Fisheries (2017): <i>Fish for the Future: Action in the Northern Basin - NSW proposal for Northern Basin Toolkit measures to promote native fish health</i> measures to overcome priority fish barriers	Fisheries
		Implement the <i>Fisheries Management Act 1994</i>	Fisheries
		Implement works required under conditions related to Burrendong Dam upgrade works	WaterNSW, Fisheries
		Seek funding opportunities to treat problematic existing structures e.g. Gin Gin, town weirs, privately owned weirs & regulators	WaterNSW, Councils, landholders, Fisheries, OEH, LLS

Risk	Description	Potential management strategies	Responsible organisations
Pumps & other water offtakes impacting on LTWP fish objectives	There are over 450 pump offtakes with a diameter greater than 200 mm located in the Macquarie-Castlereagh.	Implement NSW Department of Primary Industries, Fisheries (2017): <i>Fish for the Future: Action in the Northern Basin - NSW proposal for Northern Basin Toolkit measures to promote native fish health</i> measures for fish-friendly water extraction	Fisheries, LLS, Industry groups
No protection of environmental flow to connect with the Barwon River impacting on LTWP fish, connectivity & Barwon flow contribution targets.	Current unregulated WSP end of system access rules do not ensure a minimum flow rate from the Macquarie/Castlereagh River System into the Barwon River. Additional end of system flows that could be provided by HEW are not protected from extraction. Actual contribution of Macquarie/Castlereagh River System flows to Barwon River cannot currently be accurately determined & managed.	Consider mechanisms to protect Held environmental water through unregulated areas downstream of the regulated system This could be done to meet in-system or connection outcomes.	DOI-W
		Improve gauging of flows & providing event-based management for events in unregulated areas where HEW will be used.	DOI-W, WaterNSW
		Upgrade the capability of end of system gauges, in combination with event-based metering of unregulated water users, to enable measurement of actual end of system flows & contributions to the Barwon River. See Table 19.	DOI-W, WaterNSW



Figure 13 Golden perch. Photo: Gunther Schmida

5.3 Dealing with the risk of climate change

Climate change is a key long-term risk to river, wetland and floodplain health. Modelling indicates it will exacerbate the natural seasonal variability that exists in NSW, making it more difficult to manage our landscapes and ecosystems and the human activities that depend on them.

According to the NARCLiM model⁴⁰ (scenario 2), the changes shown in Table 14 are predicted across the Central West and Orana district by 2030 and 2070.

There are uncertainties with these climate change predictions, and the predicted changes will not occur in isolation. Rather, the predicted changes will occur alongside other changes owing to water resource development, land use, and environmental water management. The potential impact of these changes on the environmental assets of the Macquarie-Castlereagh WRPA has not been specifically assessed.

Identifying climate-related risks to water management

Environmental water management has been occurring in the Macquarie-Castlereagh WRPA since the 1980s. The climate has been variable during this time, with the region experiencing extreme droughts and floods. Environmental water and river managers have become experienced in dealing with highly variable conditions, using risk-based, adaptive management practices established over time based on real-world experience.

It will take many years to confirm that there has been a genuine change in the climate in the catchment, and in the meantime the range of management strategies already in place to deal with variable weather are likely to be relevant for managing the health of the catchment in a changing climate.

Table 13 Projected change in rainfall patterns and weather in the Central West and Orana district from 2020 to 2079 due to climate change NARCLiM (nd)

Potential change due to climate change	Description of projected change		NARCLiM projection (scenario 2)	
			2020–39	2060–79
Change in rainfall	By 2030 there will be little change in total annual rainfall. Rainfall will increase across the region during autumn & spring. Rainfall will decrease across the region during summer & winter.	Summer	-1.1%	+13.2%
		Autumn	+14.7%	+13.5%
		Winter	-4.2%	+5.4%
		Spring	+7.6%	-5.8%
Change in average temperature	Mean temperatures are projected to rise by 0.7°C by 2030. The greatest increase would be during summer & spring.	Summer	+0.95C	+2.44C
		Autumn	+0.65C	+2.04C
		Winter	+0.40C	+1.65C
		Spring	+0.80C	+2.30C
Change in number of hot days (max. temp. >35C)	Hots days are projected to increase across the region by an average of 7 days per year by 2030.	Annual	+9.1	+27.0

⁴⁰ The NARCLiM (n.d.) projections have been generated from four global climate models (GCMs) dynamically downscaled by three regional climate models (RCMs).

Potential change due to climate change	Description of projected change		NARCLiM projection (scenario 2)	
			2020–39	2060–79
Change in number of cold nights (min. temp. <2C)	Cold nights are projected to decrease by an average of 9 days per year by 2030. Changes in cold nights can have considerable impacts on native ecosystems.	Annual	-7.7	-22.5
Bushfires changes in number of days a year FFDI>50 ⁴¹	Overall, severe fire weather is projected to increase (slightly) across the region by 2030. However, increased severe fire weather is expected in the north-west part of the region during spring (the prescribed burning season) & summer (peak fire risk season). Conversely, declines in severe fire weather are expected in autumn due to increases in rainfall.	Annual	+0.5	+1.3
Hillslope erosion	Increased are predicted in average hillslope erosion rates due to changes.	Mean percent change	4.6%	20.0%
Biodiversity	Species composition will likely be changed by the suite of changes described above.			

⁴¹ Forest Fire Danger Index (FFDI) is used in NSW to quantify fire weather. The FFDI combines observations of temperature, humidity and wind speed. Fire weather is classified as severe when the FFDI is above 50.

6. Water management under different water availability scenarios

6.1 Prioritisation of ecological objectives and watering in regulated river reaches

Environmental water managers and environmental water advisory groups consider a range of factors when determining how discretionary water for the environment should be managed. Key considerations include:

- the current condition of the plants and animals, the recent history connectivity of river channels to their floodplain systems, rainfall history and predictions, and
- water availability for the range of environmental water sources.

Planning for the management of flow-dependent environmental assets amid this variability means that plans must be adaptive. They need to accommodate watering activities that range from maximising environmental outcomes from flow events when water is abundant, to managing water to maintain drought refuges when resources become scarce. Appropriate compliance activities to prevent unauthorised extractions is paramount to the success of any water management strategy's ability to contribute to environmental outcomes.

Sections 6.1.1 to 6.1.4 set out a framework to help inform annual water management decisions in river reaches which are regulated or affected by regulated water. This information is presented in terms of a water resource availability scenario (RAS) as proposed by MDBA (2012b). Each RAS is described below in two tables that include:

- in the upper tables, the broad priorities that are likely to apply to management under the particular water availability scenario along with management strategies for achieving these priorities - a 'toolkit' of opportunities to consider.
- In the lower tables (Table 14 to Table 17) the priority LTWP objectives identified for each scenario. These tables also outline the flow types which would be required to support those priority objectives.

Note the wording of the LTWP objectives has been adjusted to highlight the most relevant aspect of the objective under the scenario. For example, an LTWP objective that over 20 years seeks 'improvement' may only seek to 'maintain' under a dry scenario. Some of the wording of the objectives has been summarised for better presentation. The full objectives can be found in Section 0.

More information about RASs and how they are defined is provided in Appendix B.

6.1.1 Water RAS: Very dry - Protect

Very dry	Broad water management priorities	Key management strategies for consideration
	Avoid critical loss of species, communities & ecosystems Maintain refuges Avoid irretrievable damage or catastrophic events Avoid unnaturally prolonged dry periods Support targeted longitudinal connectivity within catchment for functional processes & a range of flora & fauna	Focus on limiting exceedance of maximum inter-flow periods, through the following strategies. Allow dry down consistent with historical wetting-drying cycles Sustain water levels & water quality in key in-channel pools, instream habitat & core wetland areas ⁴² Provide targeted low flows to allow limited fish movement Prevent two consecutive years of extreme dry to core wetland areas When available, use supplementary allocations to maximise the benefit of larger tributary flow events to meet EWRs or floodplain & lower-inundation frequency wetland assets. If a critical water shortage or similar critical incident restricts the use of water for the environment, then OEH, as part of the Critical Water Advisory Panel, will work to minimise exceedances of maximum inter-flow periods for core areas.

Table 14 Priority LTWP objectives and flow components in a very dry RAS

Priority LTWP objectives	Flow components							
	Cease-to-flow	Very Low Flow	Baseflow	Small Fresh	Large Fresh	Small Overbank /Wetland	Medium/Large Overbank /Wetland	
NF1: No loss of native fish species								
NV1: Maintain non-woody vegetation communities occurring within channels								
NV2: Maintain non-woody vegetation communities occurring in core wetlands								
WB1: Maintain waterbird species								
WB2: Maintain waterbird abundance								
WB5: Maintain waterbird habitats								
EF1: Provide & protect refugia								
EF2: Maintain quality instream & wetland habitat								
OS1: Maintain flow-dependent frogs								

⁴² Alternative watering actions such as pumping may be required to support floodplain habitats to prevent threatened fish species from drying out.

6.1.2 Water RAS: Dry - Maintain

	Broad water management priorities	Key management strategies for consideration
Dry	<p>Support the survival & viability of threatened species & communities</p> <p>Maintain refuges</p> <p>Maintain environmental assets & ecosystem functions</p> <p>Avoid unnaturally prolonged dry periods</p> <p>Support longitudinal connectivity for functional processes & a range of flora & fauna</p>	<p>Avoid exceedance of maximum inter-flow periods & provide events which have recently had lower than ideal frequency, through the following strategies.</p> <ul style="list-style-type: none"> Sustain water levels & water quality in key in-channel pools, instream habitat & core wetland areas⁴² Provide low flows to allow limited fish movement Provide freshes to channels & flows to core wetland areas at ecologically relevant times When available, use supplementary allocations to augment the above.

Table 15 Priority LTWP objectives and flow components in a dry RAS

Priority LTWP objective	Flow components								
	Cease-to-flow	Very Low Flow	Baseflow	Small Fresh	Large Fresh	Small Overbank /Wetland	Medium/Large Overbank /Wetland		
NF1: No loss of native fish species									
NF2: Maintain short to moderate-lived generalist native fish									
NF3: Maintain short to moderate-lived floodplain specialist native fish									
NF4: Maintain moderate to long-lived flow pulse specialist native fish									
NF5: Maintain moderate to long-lived riverine specialist native fish									
NF6: Maintain mature (harvestable sized) golden perch & Murray Cod									
NV1: Maintain non-woody vegetation communities occurring within channels									
NV2: Maintain non-woody vegetation communities occurring in wetlands									
WB1: Maintain waterbird species									
WB2: Maintain waterbird abundance									
WB5: Maintain waterbird habitats									
EF1: Provide & protect refugia									
EF2: Maintain instream & wetland habitat									
EF3a: Provide movement & dispersal opportunities within catchments									
EF4: Support instream & floodplain productivity.									
OS1: Maintain flow-dependent frogs									

Core & pink zone of Marshes

6.1.3 Water RAS: Moderate – Maintain and Recover

	Broad water management priorities	Key management strategies for consideration
Moderate	Enable growth, reproduction & small-scale recruitment for a diverse range of flora & fauna	Seek to meet ideal event frequencies, prioritising EWRs that have recently had long inter-flow periods or lower than ideal frequency, through the following strategies. Provide freshes at ecologically relevant times
	Promote low-lying floodplain-river connectivity	Improve condition of key off-channel waterholes (alternative watering actions may be required to support floodplain habitats to prevent threatened fish species from drying out)
	Support medium flow river & floodplain functional processes	Build on natural events where possible to provide wetland & floodplain inundation at ecologically relevant times ⁴³
	Support longitudinal connectivity within & between catchments for functional processes & a range of flora & fauna	Provide, when watering the Marshes, flow connectivity to the Barwon River ⁴⁴
	Support low flow lateral connectivity & end of system flows	Use supplementary allocations to provide additional longitudinal & lateral connectivity during supplementary flow events
	Set aside water for use in drier years	Consider carrying over water to support water use in drier years ⁴⁵

Table 16 Priority objectives and flow components in a moderate RAS

Priority LTWP objective	Flow components								
	Cease-to-flow	Very Low Flow	Baseflow	Small Fresh	Large Fresh	Small Overbank /Wetland	Medium Overbank /Wetland	Large Overbank /Wetland	
NF1: No loss of native fish species									
NF2: Improve short to moderate-lived generalist native fish									
NF3: Improve moderate-lived floodplain specialist native fish									
NF4: Improve moderate to long-lived flow pulse specialist native fish									
NF5: Improve moderate to long-lived riverine specialist native fish									
NF6: Increase mature (harvestable sized) golden perch & Murray Cod									
NF7: Expand key short to moderate-lived floodplain specialist native fish into new areas									
NF8: Expand key moderate to long-lived riverine specialist native fish into new areas									

⁴³ Includes extending duration of flows to support waterbird colonies if they establish and need intervention

⁴⁴ Connectivity depends on suitable arrangements to protect these flows through to the Barwon.

⁴⁵ See discussion on need for carryover and further work at Section 0 and at Table 19.

Priority LTWP objective	Flow components								
	Cease-to-flow	Very Low Flow	Baseflow	Small Fresh	Large Fresh	Small Overbank /Wetland	Medium Overbank /Wetland	Large Overbank /Wetland	
NF9: Expand key moderate to long-lived flow pulse specialist native fish into new areas									
NV1: Maintain non-woody vegetation communities occurring within channels									
NV2: Maintain non-woody vegetation occurring in wetlands & on floodplains									
NV3: Maintain river red gum communities closely fringing river channels									
NV4a:									
NV4b: Maintain river red gum forests & woodlands & lignum shrublands									
NV4e:									
WB1: Maintain waterbird species									
WB2: Increase waterbird abundance									
WB3: Increase opportunities for non-colonial waterbird breeding									
WB4: Increase opportunities for colonial waterbird breeding									
WB5: Maintain waterbird habitats									
EF1: Provide & protect refugia									
EF2: Create instream, floodplain & wetland habitat									
EF3a: Provide movement & dispersal opportunities within catchments									
EF3b: Provide movement & dispersal opportunities between catchments									
EF4: Support instream & floodplain wetland productivity									
EF5: Support nutrient, carbon & sediment transport & exchange									
EF6: Support groundwater conditions to sustain groundwater-dependent biota									
EF7: Contribution of flows to the Barwon									
OS1: Maintain flow-dependent frogs									
OS2: Breeding for flow-dependent frogs									

6.1.4 Water RAS: Wet – Maintain and Improve

	Broad water management priorities	Key management strategies for consideration
Wet	<p>Enable growth, reproduction & large-scale recruitment for a diverse range of flora & fauna</p> <p>Support longitudinal connectivity within & between catchments for functional processes & a range of flora & fauna</p> <p>Support high flow lateral connectivity & end of system flows</p> <p>Set aside water for use in drier years</p>	<p>Provide events at ideal frequencies.</p> <p>Balance water use with the need to set aside water for dry times – carry over water to support water use in drier years⁴⁵.</p> <p>Where possible, build on natural events to provide wetland & floodplain inundation at ecologically relevant times⁴³ & connectivity to the Barwon⁴⁴</p> <p>Protect naturally occurring floodplain wetland inundating events & high flow connectivity with the Barwon River.⁴⁶</p> <p>Use supplementary allocations to provide additional longitudinal & lateral connectivity during supplementary flow events</p>

Table 17 Priority LTWP objectives and flow components in a wet RAS

Priority LTWP objective	Flow components								
	Cease-to-flow	Very Low Flow	Baseflow	Small Fresh	Large Fresh	Small Overbank /Wetland	Medium Overbank /Wetland	Large Overbank /Wetland	
NF1: No loss of native fish species									
NF2: Improve short to moderate-lived generalist native fish									
NF3: Improve moderate-lived floodplain specialist native fish									
NF4: Improve moderate to long-lived flow pulse specialist native fish									
NF5: Improve moderate to long-lived riverine specialist native fish									
NF6: Increase mature (harvestable sized) golden perch & Murray Cod									
NF7: Expand key short to moderate-lived floodplain specialist native fish into new areas									
NF8: Expand key moderate to long-lived riverine specialist native fish into new areas									
NF9: Expand key moderate to long-lived flow pulse specialist native fish into new areas									
NV1: Increase non-woody vegetation communities occurring within channels									

⁴⁶ Reliant on effective policy on floodplain harvesting, no increase in regulation and other measures of policy and compliance as outlined for unregulated river reaches at Table 18

Priority LTWP objective		Flow components							
		Cease-to-flow	Very Low Flow	Baseflow	Small Fresh	Large Fresh	Small Overbank /Wetland	Medium Overbank /Wetland	Large Overbank /Wetland
NV2: Maintain non-woody vegetation occurring in wetlands & on floodplains									
NV3: Maintain river red gum communities closely fringing river channels									
NV4a:	Maintain extent & maintain or increase the condition of river red gum, black box, coolibah & lignum								
NV4b:									
NV4c:									
NV4d:									
NV4e:									
WB1: Maintain waterbird species									
WB2: Increase waterbird abundance									
WB3: Increase opportunities for non-colonial waterbird breeding									
WB4: Increase opportunities for colonial waterbird breeding									
WB5: Improve waterbird habitats									
EF1: Protect refugia									
EF2: Create instream, floodplain & wetland habitat									
EF3a: Provide movement & dispersal opportunities within catchments									
EF3b: Provide movement & dispersal opportunities between catchments									
EF4: Support instream & floodplain wetland productivity									
EF5: Support nutrient, carbon & sediment transport & exchange									
EF6: Support groundwater conditions to sustain groundwater-dependent biota									
EF7: Contribution of flows to the Barwon									
OS1: Maintain flow-dependent frogs									
OS2: Breeding for flow-dependent frogs									

6.2 Protection of ecologically important flow components in unregulated river reaches

In unregulated areas, where water cannot be delivered through a regulating structure, the primary means of protecting environmentally important flows is through pumping access rules in the water sharing plans (WSPs). Table 18 sets out potential management strategies that could be implemented in the WSPs to ensure important flows are protected.

Table 18 Potential management strategies for meeting LTWP objectives in unregulated river reaches of the Macquarie Castlereagh WSPA

Flow component	Potential management strategies in reaches when issues arise	Most relevant weather scenarios
Cease-to-flow	Maintain the pools policy (no pool drawdown permitted). Consider reviewing cease-to-pump rules to reduce the length of cease-to-flow periods. Consider implementing a first flush rule to ensure cease-to-flow periods are broken at ecologically relevant times & with events of sufficient magnitude to avoid adverse water quality incidents.	Very dry Dry Moderate ⁴⁷
Low flows & baseflows	Consider rostering landholder water access during low flow months. Consider reviewing cease-to-pump rules to better protect low flows & baseflows.	Very Dry Dry Moderate ⁴⁷
Freshes	Consider implementing total &/or individual daily extraction limits (IDELS & TDELS) during ecologically important periods. ^{48,49}	Very dry Dry Moderate
Entire flow regime, including overbank & wetland inundating flows	Consider targeted water access licence purchases from willing sellers where flows are acutely impacted. Ensure compliance with water access licence conditions ⁴⁹ Protect water for the environment that originates from held water entitlements & the EWA. ^{50, 49, 51} Maintain 'no trade into water source' rules in the WSPs. Use trade restrictions to protect sensitive water sources from greater impact. Consider restrictions to take in water sources bordering the Barwon River when embargoes on take exist in the Barwon River. Monitor for changes in water demand & review access rules if current usage is high or if the pattern of use changes.	All weather scenarios

⁴⁷ Planning units lower in the WSPA may be impacted under moderate flows

⁴⁸ Individual daily extraction limits or total daily extraction limits for a particular flow class may be considered to reduce extraction pressure on ecologically important flow components

⁴⁹ May require improved water metering and gauging in certain areas

⁵⁰ Requires adequate compliance measures

⁵¹ In line with Matthews Reports (2017a,b) recommendations for protecting environmental flows

7. Going forward

7.1 Potential further actions for water management.

Table 19 outlines a number of significant investment priorities that could improve water management.

7.2 Cooperative water use

Cooperative river operations

The BWS notes that all water in the Murray–Darling Basin, including natural events and consumptive water, has the potential to contribute to improving the ecological condition of rivers, wetlands and floodplains (MDBA 2014a). Making the best use of all water is a key strategy to achieve the objectives in this LTWP. In some cases, river operating practices need to be revised to provide the operators with a mandate to manage rivers so that environmental outcomes can be achieved. The risks and constraints to achieving EWRs (Table 11) described in this LTWP identifies some river management practices that are currently limiting the ability to achieve ecological objectives. The LTWP identifies the following strategies to improve the benefit of all water in the system:

- Investigate options for the delivery of irrigation orders to more closely mimic natural flow events.
- Establish better channel sharing arrangements by permitting environmental water to build on consumptive or stock and domestic deliveries to achieve better flow regimes for the environment.
- Optimise water releases from Windamere and Burrendong Dam to mimic natural rates of fall.

Coordinating deliveries of held environmental water with consumptive deliveries can help to achieve greater flow volumes from the smarter use of all water. Such arrangements could enable larger in-channel and, where permitted, overbank flows that would not be possible with designated environmental water alone.

Similarly, controlled river flows through the system for consumptive deliveries can also meet the volumes described in many EWRs⁵², without any contribution of environmental water. One of the primary recommendations of this LTWP is to investigate the potential to optimise these outcomes, by supporting collaboration between OEH and WaterNSW to assist in shaping consumptive deliveries to meet environmental water requirements and strike a balance between operational efficiency and ecological objectives.

Cooperative environmental water management

Managing water for the environment at the catchment scale requires cooperation between stakeholders. Such cooperative arrangements support better environmental outcomes.

Water for the environment in NSW is managed cooperatively by three government agencies: OEH, CEWO and DOI-W. Together these agencies manage NSW and Commonwealth held environmental water portfolios (OEH and CEWO), discretionary planned environmental water (OEH) and the WSP rules that provide environmental benefits throughout the system (DOI-W).

⁵² Consumptive deliveries can replicate the hydrologic characteristics, however other benefits such as nutrient transport may be removed when the water is extracted

7.3 Complementary actions

Cooperative natural resources management for environmental outcomes

To achieve the watering required to support the environmental objectives, it is necessary to ensure that any priority environmental assets and functions on private land can be accessed for management. This includes arrangements with landholders that allow for priority assets on private land to be inundated with the required timing, frequency and duration. Access to these assets to evaluate how they are responding to management over time is also vital for the full implementation of the LTWP in these areas.

Complementary management of flow-dependent environmental assets is vital to the success of this LTWP. Degradation of assets through poor land management practices and inadequate legislative protection may undermine the benefits of environmental water management. Cooperative arrangements between government agencies such as LLS, private industry groups, individual landholders and community groups that ensure adequate stewardship of environmental assets are essential to the success of this LTWP.

Cooperative investment opportunities

A few significant investment priorities have been identified in the Macquarie-Castlereagh (Table 19). Identification of funding opportunities and subsequent implementation of projects to address these priorities would contribute significantly to the environmental outcomes identified in this plan.

Through the life of the plan, OEH will seek opportunities to build links and partnerships to support implementation of projects that will contribute to the ecological objectives of the LTWP.



Figure 14 Burrendong Dam offtake tower with cold water curtain. Photo: Tim Hosking/OEH

Table 19 Recommended further investment and projects to improve environmental water outcomes in the Macquarie-Castlereagh WRP

Investment opportunity	Description	Potential project partners	Relevant LTWP section
Carryover scenario modelling	Use long-term hydrological models to further inform the risk-based decisions on carryover for HEW & discretionary EWA, including benefits & opportunity costs of different carryover strategies.	DOI-W, CEWO, WaterNSW	0
Improve flow & asset information in the distributary (effluent) creeks, including Marra Creek.	Distributary creeks such as Marra & Crooked creeks have variable channel morphologies & often multiple ungauged input flow paths during higher flows. Observations during actual flow events & further channel cross sections & rating estimations could help in improving flow threshold estimates for baseflows, freshes & overbanks & in determining which flows, if any, are feasible for environmental water managers to consider. Further information on environmental values & assets would also assist to inform decisions.	DOI-W, CEWO, WaterNSW	0
Determine the feasibility of delivering water to the Talga Wetland/Overflow of the Lower Crooked Creek	This may involve a trial of controlled watering using combined unregulated flows & managed environmental water. The flow would need to be carefully monitored to see the level of inundation in the Talga Wetland area & to determine constraints to flow including structures. Prior to any water use, ideally, the creek should be surveyed for constraints & to determine if the channel capacity in the Talga Wetland area is greater than 100 ML/day capacity (Sri Sritharan, WaterNSW, pers. comm. August 2018) to avoid any ineffective use of water. Other flowpaths, such as one outlined in Torrible et al. (2011) should also be considered, though benefits should be balanced against costs & potential on-route losses. This water use would also need to be balanced against other annual priorities.	DOI-W, CEWO, WaterNSW	0
Investigate options for improved flow variability in distributary/effluent creeks	Several creeks such as Gunningbar & Duck creeks have constant baseflows but poor flow variability & less freshes. Barma et al. (2011) noted the in-stream ecology in these creeks was degraded due to poor flow variability & considered the creeks a poor target for environmental flows until a more variable low flow regime can be implemented. Methods to ensure users have consistent water supply, such as on-property storage may allow some greater flow variability, including periods of natural drying as recommended by Torrible et al. (2011), & potential use of the water savings for freshes. Any reductions in baseflows, & particularly any periods of drying down, would have to be considered with regard to fish communities & refuge pools which are likely to be shallow due to sedimentation.	DOI-W, CEWO, WaterNSW	0
Increase the number of gauges in key locations	Key end of system gauges. Current end-of-system gauges are upstream of unregulated licenced extraction points. It is therefore not possible to accurately measure flows to the Barwon. These flow values are important to determine if sufficient connectivity has been	DOI-W, WaterNSW	5.1

Investment opportunity	Description	Potential project partners	Relevant LTWP section
	<p>achieved & flow contributions to the Barwon have been provided. They would also support active management of events in these reaches.</p> <p>Key unregulated watercourses. In unregulated systems where no gauges exist, commence-to-pump rules are often unable to specify flow rates, which is important to protect critical low & baseflows. Gauges should be installed to allow better flow monitoring in these locations.</p>		
Increase outlet valve capacity increase at Burrendong Dam	The maximum valve capacity at Burrendong Dam at Full Supply Level is 8500 ML/d & less at low dam levels. Increased outlet capacity will improve wetland watering opportunities & reduce potential conflict between the environment & consumptive water users.	DOI-W, Water NSW	5.1
Raise flow constraint on the Cudgegong River at Rocky Waterhole Bridge	Cudgegong River flows are restricted to around 1,500 ML/day by the Rocky Waterhole Bridge. Support proposed upgrade of Cudgegong River crossing to allow higher flows.	DOI-W, WaterNSW	5.1
Investigate options for relaxing constraints to delivery of environmental flows to the Macquarie Marshes	<p>Environmental flows to the Marshes via Marebone Weir are currently constrained to 3,200-3,500 ML/d⁵³ to avoid breakouts from the river, particularly towards Crooked Creek. Flow rates between 3,200 ML/day & 4,000 ML/day are valuable to achieving floodplain connection in some areas, such as the southern part of the Eastern Marshes through 'the Jungle' (Back Swamp) & other flow paths (D Love, OEH, pers. comm. August 2018).</p> <p>A smaller version of the constraints management strategy projects (MDBA 2014b) in other valleys, such as the Gwydir, would benefit environmental water management in the Marshes.⁵⁴</p>	DOI-W, WaterNSW	5.1
Investigate channel works to reinstate appropriate channel profile in selected streams in the Southern Macquarie Marshes	Continue to investigate channel works in the Southern Marshes to stabilise & restore channel profiles, allowing continued & enhanced connection with wetlands & the floodplain. Priorities for works are identified in the Southern Macquarie Marshes Scoping Study (Ralph et al. 2013).	DOI-W, WaterNSW, NPWS	5.1

⁵³ Note EWA translucent flows are able not as restricted and are able to be delivered up to 4,000 ML/day.

⁵⁴The MDBA (2013) preliminary overview of constraints in the Murray-Darling Basin did not identify a first order constraint in the Macquarie-Castlereagh and consequently the catchment was not prioritised for funding under the Constraints Management Strategy (MDBA 2014b). However, the preliminary overview did not investigate important local issues such as access ways and cropping in flow paths to the Marshes. Flows to the East Marshes are particularly affected by this.

Investment opportunity	Description	Potential project partners	Relevant LTWP section
Protect flows to the Barwon at critical times	Work with DOI-W & others to investigate ways to limit access to end of system flows in the Macquarie-Castlereagh system when low flows have restricted access in the Barwon River downstream of the relevant Macquarie/Castlereagh WRPA waterway confluences.	DOI-W, WaterNSW	5.2
Provide incentives to improve management of wetlands on private land	<p>The protection of native vegetation requires good knowledge, the cooperation of various stakeholders, & multiple different projects, which could include:</p> <ul style="list-style-type: none"> • habitat mapping to identify riparian & aquatic habitat condition & prioritise reaches for management actions in partnership with LLS & landholders, to develop formal agreements & unified strategies • implement grazing strategies required to protect & restore wetland vegetation, bank stability & adequate water quality in collaboration with LLS & landholders • provide incentives to landholders to improve management of wetlands on private land. 	OEH, LLS, NPWS, CEWO, Fisheries, DPI, Landholders, Biodiversity Conservation Trust	5.2
Formalise channel sharing arrangements	Develop formal supply sharing arrangements to provide a mechanism for managing the delivery of water when demands regularly exceed channel sharing capacity.	OEH, DOI-W, Water NSW, CAG	
Implementation of a native fish restoration project	<p>To assist in improving the aquatic habitat that supports native fish there is an opportunity to implement various instream management activities, including:</p> <ul style="list-style-type: none"> • assessing & addressing priority barriers to fish passage in the WRPA • estimating the flow rates required to drown out barriers and provide connectivity through important river reaches (for example, PUs that connect to the Barwon) • the implementation of pump screening methods to prevent entrainment of native fish, larvae & eggs • works to achieve instream habitat improvement including re-snagging & aquatic revegetation • instream habitat mapping to help identify high-risk & priority refuge areas • implementation of the carp management strategy (see NSW I&I 2010) and National Carp Control Plan (FRDC, in prep). • reintroduction, translocation & stocking of threatened fish species in key locations. <p>Note some aspects of this project may have funding under <i>Fish for the Future: Action in the Northern Basin—NSW proposal for Northern Basin Toolkit measures to promote native fish health</i> (Fisheries, 2017)</p>	Fisheries, LLS	5.2

Investment opportunity	Description	Potential project partners	Relevant LTWP section
Assessment of first flush requirements in select PUs	Using information on high-risk & priority refuge pools determined in the native fish restoration project (above), estimate the magnitude of flow required to flush these pools following a CTF event which would reduce the risk of adverse water quality events. This may require water quality monitoring and additional river gauges.	OEH, Fisheries, DOI-W	
Improve understanding of climate change impacts, thresholds & adaptive management strategies	Potential changes in weather, bushfires & erosion due to climate change are outlined in Table 13. This task involves assessing the potential impact of these changes on the environmental assets of the Macquarie-Castlereagh WRPA, identifying management adaptations to respond to change & developing indicators to trigger responses.	Fisheries, NPWS, DOI-W.	

7.4 Measuring progress

Monitoring, evaluating and reporting (MER) to support adaptive management are integral to informing planning and operational decisions. Monitoring how water moves through the system and how the environment responds informs ongoing improvements to water management. This information also assists in informing revisions of this LTWP every five years.

Monitoring and evaluating environmental water management in the Macquarie-Castlereagh WRPAs draws on contributions from Australian and NSW Government agencies, universities, other research organisations, non-government organisations, individuals and land managers.

The MER program provides a unified approach to delivering Basin Plan and NSW evaluation and reporting requirements. The NSW-wide MER program consists of:

- a NSW MER Framework that describes the principles, types of monitoring, alignment across NSW agencies efforts, knowledge gaps, externalities and constraints, and relationships to the BWS and Basin Plan. It also describes how existing knowledge and programs are used to create a cost-effective and coordinated program
- the OEH-specific parts, called the *Healthy Inland Wetlands Environmental Water Program* that describes the approach to developing LTWP MER objectives, evaluation of management actions, and reporting
- customised MER Plans that summarise the proposed integrated MER activities for each WRPAs
- monitoring *Methods Manuals* that describe methods for each monitoring theme (e.g. fish, hydrology, vegetation, water quality, macroinvertebrates, waterbirds) considered in broader NSW water monitoring. These manuals, when developed, will contain information relating to survey, data handling and analysis techniques, conceptual models and cooperative research arrangements.

The NSW MER Framework, which includes NSW Fisheries Basin Plan Environmental Outcomes Monitoring program and OEH's MER program, provides the structure within which various NSW-led monitoring activities are brought together for:

- tracking progress towards stated LTWP and WSP outcomes
- improved decision making for environmental water planning and operations (supporting adaptive management).

To do this, the 2018 NSW MER Framework, aim to:

- evaluate progress towards achieving outcomes defined within LTWPs and WSPs
- extend, augment and respect current and historical monitoring
- address information and monitoring gaps or short-falls
- provide high-quality, scientifically-robust information to support both continual improvement of operations and a growing information base for wetland and river conservation generally
- collaborate with water delivery partners (particularly the CEWO), DoI-W, wetland managers, other agencies and researchers to value-add to monitoring outcomes and minimise duplication
- provide information that supports community engagement and improved reporting of environmental water outcomes which will increase government and community confidence and awareness of environmental water management

- streamline reporting requirements under WRPs, LTWPs, Schedule 12 of the Basin Plan and the National Partnership Agreement.

The detail of the monitoring to be undertaken under the OEH MER program is being finalised and is dependent on the level of available funding.

Monitoring progress reports are made available following each watering year.

7.5 Review and update

This LTWP brings together the best available information from a range of community, traditional and scientific sources. To ensure the information, objectives and supporting material remains relevant and up-to-date, this LTWP will be reviewed and updated no later than five years after it is implemented, being 2024. Additional reviews of all, or part, of this LTWP may also be triggered by:

- accreditation or amendment to the WSP or WRP for the Macquarie-Castlereagh WRPA
- revision of the BWS that materially affects this LTWP
- a sustainable diversion limit (SDL) adjustment
- significant new information arising from evaluating responses to environmental watering
- significant new information about the ecology of the Macquarie-Castlereagh WRPA that is relevant to environmental water management
- improved understanding of the effects of climate change and its impacts on the Macquarie-Castlereagh WRPA
- significant changes to the river operating environment or the removal of constraints that affect watering strategies
- material changes to river and wetland health, not considered within this LTWP.

References

- Adams SB & Warren LM 2005, Recolonization by warmwater fishes and crayfishes after severe drought in upper Coastal Plain hill streams, *Transactions of the American Fisheries Society*, 134: 1173–1192.
- Alluvium 2010, *Key ecosystem functions and their environmental water requirements*, Report by Alluvium to the Murray-Darling Basin Authority, Canberra.
- Amtstaetter F, O'Connor J & Pickworth A 2016, Environmental flow releases trigger spawning migrations by Australian grayling *Prototroctes maraena*, a threatened, diadromous fish, *Aquatic Conservation: Marine and Freshwater Ecosystems*, 26: 35–43.
- Arthington A, Olden J, Balcombe S and Thoms M, 2010, Multi-scale environmental factors explain fish losses and refuge quality in drying waterholes of Cooper Creek, an Australian arid-zone river, *Marine and Freshwater Research*, 61(8).
- Baldwin DS, Colloff MJ, Mitrovic SM, Bond NR & Wolfenden B 2016, Restoring dissolved organic carbon subsidies from floodplains to lowland river food webs: a role for environmental flows? *Marine and Freshwater Research*, 67: 1387–1399.
- Barma Water Resources in association with IRPEC Pty Ltd and Paul Wettin 2011, *Environmental Water Delivery: Macquarie River*, Prepared for Commonwealth Environmental Water, Department of Sustainability, Environment, Water, Population and Communities, Canberra.
- Boulton AJ 2003, Parallels and contrasts in the effects of drought on stream macroinvertebrate assemblages, *Freshwater Biology*, 48(7): 1173–1185.
- Bowen S & Simpson SL 2010, *Changes in the extent and condition of the vegetation communities of the Macquarie Marshes floodplain 1991–2008: Final Report to the NSW Wetland Recovery Program*. Rivers and Wetlands Unit, Department of Environment Climate Change and Water, NSW, Sydney.
- Brandis K 2010, *Colonial waterbird breeding in Australia: wetlands, water requirements and environmental flows*, PhD Thesis. Australian Wetlands and Rivers Centre, School of Biological, Earth and Environmental Sciences, University of New South Wales. Sydney.
- Brandis KJ & Bino G 2016, Habitat use by waterbirds in and adjacent to the Murray-Darling Basin, MD3360, The University of New South Wales, Sydney.
- Casanova MT 2015, *Review of water requirements for key floodplain vegetation for the northern basin: literature review and expert knowledge assessment*, Report to the Murray-Darling Basin Authority, Charophyte Services, Lake Bolac.
- DECCW 2010, *NSW Climate Impact Profile – the impacts of climate change on the biophysical environment of New South Wales*, NSW Department of Environment, Climate Change and Water.
- DECCW 2011, *NSW Rivers Environmental Restoration Program, Final Report*, NSW Department of Climate Change and Water, Sydney.
- DOI-W in prep, Risk Assessment for the Macquarie-Castlereagh Water Resource Plan Area (SW11), NSW Department of Industry - Water.
- DOI-W in prep, Water Quality Salinity Management Plan for the Macquarie-Castlereagh Water Resource Plan Area. NSW Department of Industry - Water.

DPI 2015, *Fish and Flows in the Northern Basin: responses of fish to changes in flow in the Northern Murray-Darling Basin – Valley Scale Report*, Final report prepared for the Murray-Darling Basin Authority, NSW Department of Primary Industries, Tamworth.

DPI 2017, *Fish for the Future: Action in the Northern Basin - NSW proposal for Northern Basin toolkit measures to promote native fish health*, NSW Department of Primary Industries.

Fairfull S & Witheridge G 2003, *Why do Fish Need to Cross the Road? Fish Passage Requirements for Waterway Crossings*, NSW Fisheries, Cronulla, 16 pp.

FRDC (in prep), *National Carp Control Plan*, Fisheries Research and Development Corporation, <www.carp.gov.au>

Gippel CJ 2013, *Assessment of the hydraulics of the Little Murray River Weir Pool under alternative operating scenarios*, Fluvial Systems Pty Ltd, Stockton. Goulburn-Murray Water, Shepparton, January.

Holmes S, Speirs S, Berney P & Rose H 2009, *Guidelines for grazing in the Gwydir Wetlands and Macquarie Marshes*, A product of the NSW Department of Primary Industries' Best Management Practice for Grazing in Wetlands project.
<www.dpi.nsw.gov.au/agriculture/pastures-and-rangelands/establishment-mgmt/grazing-management2/guidelines-gwydir-macquarie>

Humphries P & King AJ 2004, Drifting fish larvae in Murray-Darling Basin rivers: composition, spatial and temporal patterns and distance drifted, In *Downstream Movement of Fish in the Murray – Darling Basin. Statement, Recommendations and Supporting Papers from a Workshop held in Canberra 3–4 June 2003* (eds M. Lintermans and B. Phillips). pp. 51–58, Murray – Darling Basin Commission, Canberra, Australia.

Kingsford RT 1999, Aerial survey of waterbirds on wetlands as a measure of river and floodplain health, *Freshwater Biology*, 41, 425-438

Kingsford RT & Auld KM 2005, Waterbird breeding and environmental flow management in the Macquarie Marshes, Arid Australia. *River Research and Applications* 21:187-200.

Langhans S & Tockner K 2006, The role of timing, duration, and frequency of inundation in controlling leaf litter decomposition in a river-floodplain ecosystem (Tagliamento, northeastern Italy), *Oecologia*, 147(3): 501-509.

Matthews K 2017a, *Independent investigation into NSW water management and compliance – interim report*, Report to the NSW Department of Primary Industries, September 2017.

Matthews K 2017b, *Independent investigation into NSW water management and compliance – final report*, Report to the NSW Department of Primary Industries, December 2017.

MDB Ministerial Council 2015, *Basin Salinity Management 2030*, Murray–Darling Basin Ministerial Council.

MDBA 2012a, *Basin Plan*, Legal instrument prepared under subparagraph 44(2)(c)(ii) of the Water Act 2007, Murray Darling Basin Authority, Commonwealth of Australia.

MDBA 2012b, *Guidelines for the method to determine priorities for applying environmental water*, Murray Darling Basin Authority, Canberra.

MDBA 2013, *Preliminary Overview of Constraints to Environmental Water Delivery in the Murray–Darling Basin: Technical Support Document*, Murray-Darling Basin Authority, Canberra.

MDBA 2014, *Basin-wide environmental watering strategy*. Murray-Darling Basin Authority, Canberra.

MDBA n.d., *Macquarie–Castlereagh – Snapshot*. Murray-Darling Basin Authority, Canberra. Accessed 1 September 2018, <www.mdba.gov.au/discover-basin/catchments/macquarie-castlereagh>

Mora-Gomez J, Elozegi A, Mas-Marti E & Romani AM 2015, Factors controlling seasonality in leaf-litter breakdown in a Mediterranean stream, *Freshwater Science*. 34(4): 1245-1258.

NARClIM n.d., NSW and ACT Regional Climate Modelling (NARClIM) Project. A multiagency research partnership between the NSW and ACT governments and the Climate Change Research Centre at the University of NSW. Available at: <https://climatechange.environment.nsw.gov.au/Climate-projections-for-NSW>. Accessed 29 August 2018.

NSW I & I 2010, *NSW Control Plan for the noxious fish carp Cyprinus carpio*. Department of Industry and Investment NSW, www.dpi.nsw.gov.au/_data/assets/pdf_file/0010/306010/nsw-control-plan-for-carp.pdf

O'Connor J, Mallen-Cooper M & Stuart I 2015, *Performance, operation and maintenance guidelines for fishways and fish passage works*, Arthur Rylah Institute for Environmental Research Technical Report No. 262 for the Water and Catchments Group, Department of Environment, Land, Water and Planning. Arthur Rylah Institute for Environmental Research, Department of Environment, Land, Water and Planning, Heidelberg, Victoria.

Ocock J & Spencer J 2018, Results from spring frog surveys in the Macquarie Marshes (2011-17). Summary report February 2018, Unpublished report, NSW Office of Environment and Heritage.

OEH 2010, *Macquarie Marshes Adaptive Environmental Management Plan*, NSW Office of Environment and Heritage.

OEH 2012, *Macquarie Marshes Ramsar Site. Ecological character description Macquarie Marshes Nature Reserve and U-block component*, NSW Office of Environment and Heritage, Sydney.

OEH 2013, *Macquarie Marshes Ramsar site - Article 3.2 response strategy*, NSW Office of Environment and Heritage, Sydney.

OEH 2018, *Draft Floodplain Management Plan for the Macquarie Valley Floodplain 2018*, NSW Office of Environment and Heritage, Sydney.

OEH (Unpublished), Monitoring manual – OEH environmental water program – Standard methods – frogs, NSW Office of Environment and Heritage, Sydney.

Poff NL & Zimmerman JKH 2010, Ecological responses to altered flow regimes: a literature review to inform the science and management of environmental flows, *Freshwater Biology*, 55(1): 194-205.

Ralph T & Rogers K 2011, Floodplain wetlands of the Murray-Darling Basin and their freshwater biota. In K Rogers & T Ralph (Eds.), *Floodplain Wetland Biota in the Murray-Darling Basin: Water and Habitat Requirements* (pp. 1-16), Collingwood, Vic: CSIRO Publishing.

Ralph T, Barma D, Hesse P, Outhet D, Davies B and Wettin P 2013, *Southern Macquarie Marshes geomorphic restoration scoping study*, Sydney, Australia: Access Macquarie Ltd.

Roberts J & Marston F 200, *Water regime of wetland & floodplain plants in the Murray-Darling Basin: A source book of ecological knowledge*, National Water Commission, Canberra.

Roberts J & Marston F 2011, *Water regime for wetland & floodplain plants: a source book for the Murray-Darling Basin: A source book of ecological knowledge*, Canberra: CSIRO Land and Water

Rogers K & Ralph TJ 2011, *Floodplain wetland biota in the Murray-Darling Basin: water and habitat requirements*, CSIRO Publishing, Collingwood, Victoria, Australia.

Spencer J, Ocock J, Amos C, Borrell A, Suter S, Preston D, Hosking T, Humphries J, Hutton K, Berney P, Lenon E, Brookhouse N, Keyte P, Dyer J, and Lenehan J 2018, Monitoring waterbird outcomes in NSW: Summary Report 2016-17, Unpublished report, NSW Office of Environment and Heritage, Sydney.




Spencer JA 2017, Evaluation of colonial waterbird breeding in NSW Murray-Darling Basin: 2006-16, Unpublished report, NSW Office of Environment and Heritage, Sydney.

Thomas RF, Jones H & Karunaratne SB In prep, Flow scenario inundation assessment for the vegetation of the Macquarie Marshes.

Torrible L, Wettin P, Barma D, Wilson G, Hobcroft D and Ocock J 2011, *Post flood assessment and determination of environmental water requirements for Gunningbar Creek, Lower Crooked Creek, Marra Creek and the Lower Macquarie River*. Prepared by IRPEC Pty for BWR on behalf of the Australian Government Department of Sustainability, Environment, Water, Population and Communities.

Westhorpe DP, Mitrovic SM & Kobayashi Y., 2010. Limitations of lowland riverine bacterioplankton by dissolved organic carbon and inorganic nutrients. *Hydrobiologia*, 320: 292-298.

Appendix A LTWP objectives relevant to each planning unit in the Macquarie-Castlereagh WRP


			1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28			
			Regulated or affected by regulated											Predominantly unregulated																			
			Macquarie River											Macquarie river, tributaries and distributaries										Upper Bogan R & Headwaters	Castlereagh and tributaries								
Code	Ecological objective		Cud reg	Barroona to Burrendong to Barroona	Warren to Barroona to Marshes	Marsh South	Marsh North	Marsh East	Lower Macq	Marra Creek	Lower Bogan	Marthaguy	Cudgegong headwaters	Macq Headwaters	Upper Macq tributaries	Upper Macq River Channel	Bell River	Little River	W-bang, Wyl, M-vale, Geurie	Talbragar River	Coolbaggie Ck	B-water Boggy	Ewenmar & B-water Boggy	Upper Bogan R & Headwaters	C-reagh above Binnaway	Binnaway to Glandarra	Toorweannah to Coonamble	C-reagh R Glandarra to Coonamble	Lower C-reagh Tribs	C-reagh below Coonamble			
<div>Native fish</div> 	NF1	No loss of native fish species	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●			
	NF2	Distribution and abundance of short to moderate-lived generalists	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●		
	NF3	Distribution and abundance of short to moderate-lived floodplain specialists			●	●	●	●	●	●	●	●										●					●	●	●	●	●		
	NF4	Population structure for moderate to long-lived flow pulse specialists	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
	NF5	Population structure for moderate to long-lived riverine specialists	●	●	●	●	●	●		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
	NF6	A 25% increase in abundance of harvestable sized Golden Perch and Murray Cod	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
	NF7	Key short to moderate-lived floodplain specialists into new areas (in historical range)							●		●											●	●								●		
	NF8	Key moderate to long-lived riverine specialists into new areas (within historical range)		●	●									●						●	●			●									
	NF9	Key moderate to long-lived flow pulse specialists into new areas (in historical range)						●	●	●		●	●																			●	
<div>Waterbirds</div> 	WB1	Number and type of waterbird present.				●	●	●	●	●																							
	WB2	Total waterbird abundance.					●	●	●	●																							
	WB3	Breeding activity in non-colonial nesters					●	●	●	●																							
	WB4	Opportunities for colonial breeding events.					●	●	●	●																							
	WB5	Condition of waterbird habitats				●	●	●	●	●	●																						
<div>Vegetation</div> 	NV1	Non-woody vegetation communities occurring within channels	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●		●	●	●	●	●	●	●	●	●	●	
	NV2	Non-woody vegetation communities occurring in wetlands and on floodplains				●	●	●	●	●	●	●										●	●									●	
	NV3	River red gum communities closely fringing river channels	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
	NV4a	Floodplain and wetland forest, woodland and shrublands		●	●	●	●	●	●	●	●	●										●	●								●	●	
	NV4b			●	●	●	●	●	●	●	●	●	●									●	●								●	●	
	NV4c				●	●	●	●	●	●	●	●	●									●	●								●	●	
	NV4d				●	●	●	●	●	●	●	●	●									●	●								●	●	
NV4e				●	●	●	●	●	●	●	●	●									●	●								●	●		

Commented [PD1]: Word 'Upper' added here.

Commented [PD2]: Dot DELETED here – for Eweremar and Backwater Boggy – for NF7

Commented [PD3]: Dot added here – Tabragar River – for NF8

Commented [PD4]: Dot added here – for Upper Bogan for NF8

			1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
			Regulated or affected by regulated											Predominantly unregulated																
Code	Ecological objective		Macquarie River							Marra Creek	Lower Bogan	Marthaguy	Macquarie river, tributaries and distributaries										Upper Bogan R & Headwaters	Castlereagh and tributaries						
			Cud reg	Burrendong to Barroona	Barroona to Warren	Warren to Marshes	Marsh South	Marsh North	Marsh East				Lower Macq	Cudgegong headwaters	Macq Headwaters	Upper Macq tributaries	Upper Macq River Channel	Bell River	Little River	W-bang., Wyl., M-vale, Geurie	Talbragar River	Coolbaggie Ck		B-water Boggy	Ewenmar & B-water Boggy	C-reagh above Binnaway	Binnaway to Gilgandra	Binnaway to Tooraweenah	Tooraweenah to Coonamble	C-reagh R Gilgandra to Coonamble
<div>Ecosystem functions</div> <div>↑↓</div>	EF1	Provide and protect a diversity of refugia	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
	EF2	Create quality instream, floodplain and wetland habitat.	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
	EF3a	Provide movement and dispersal opportunities within catchments.	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
	EF3b	Provide movement and dispersal opportunities between catchments.	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
	EF4	Support instream and floodplain productivity.	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
	EF5	Support mobilisation and transport of sediment, carbon and nutrients.	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
	EF6	Support groundwater conditions to sustain groundwater dependent biota.	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
	EF7	Increase the contribution of flows into the Murray and Barwon-Darling tributaries.							●	●	●	●																		●
<div>Other Species (frogs)</div> <div></div>	OS1	Maintain species richness and distribution of flow-dependent frog communities					●	●	●																					
	OS2	Maintain successful breeding opportunities for flow-dependent frog species					●	●	●																					

Commented [PD1]: Word 'Upper' added here.

Appendix B RAS guidelines

In the Macquarie-Castlereagh WRPA the Environmental Water Advisory Group uses RASs to guide decision-making. The formal assessment of the RAS occurs twice during the planning process: preliminary planning (February) and detailed planning (May).

The critical information required for this assessment is water availability in accounts, allowances and the assessed condition of the environment (antecedent conditions). As set out in section 8.61 of the Basin Plan a RAS will be one of: very dry, dry, moderate, wet, or very wet.

To determine the RAS, the following steps are followed⁵⁵:

- a) determine the antecedent conditions for a given water resource plan area by (the 'X' axis of the matrix in Table 21):
 - i. selecting a representative number of water accounting periods preceding the current water year (e.g. 3–5 years)
 - ii. assessing the water received by the environment for those years
 - iii. comparing the amount in (ii) to all the historical data
 - iv. categorising the antecedent conditions as a percentile relative to all historical water years
- b) determine the surface water availability by (the 'Y' axis of the matrix in Table 20):
 - i. assessing all sources of water available for the environment for a given period
 - ii. comparing these to all the historical data
 - iii. categorising the surface water availability as a percentile relative to all historical water years
- c) for the relevant water accounting period, determine the surface water availability relative to the antecedent conditions for the water resource plan area using all of the historical climate condition data that are available (in Table 20, this is the surface water availability percentile)
- d) using the following matrix below.

Table 20 Default matrix for determining the RAS (MDBA 2012b)

Surface water availability	Antecedent conditions				
	Very dry (0–15%)	Dry (16–45%)	Medium (46–60%)	Wet (61–85%)	Very wet (86–100%)
Very low (0–15%)	Very dry	Very dry	Dry	Dry	N/A
Low (16–45%)	Very dry	Dry	Dry	Moderate	Wet
Medium (46–60%)	Dry	Dry	Moderate	Wet	Wet
High (61–85%)	Dry	Moderate	Wet	Wet	Very wet
Very high (86–100%)	N/A	Moderate	Wet	Very wet	Very wet

⁵⁵ As outlined in MDBA (2012b)